

PARASITOLOGY

LABORATORY ONLY

CDC

Dr. HEYNEMAN

PICTORIAL KEYS

**ARTHROPODS, REPTILES, BIRDS AND MAMMALS
OF PUBLIC HEALTH SIGNIFICANCE**

**U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE**

**PICTORIAL KEYS TO
ARTHROPODS, REPTILES, BIRDS AND MAMMALS
OF PUBLIC HEALTH SIGNIFICANCE**



**U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
Communicable Disease Center
Atlanta, Georgia 30333**

1966

C O N T E N T S

Introduction	1
Pictorial Keys:	
General	2
Crustacea	16
Centipedes	19
Millipedes	20
Arachnida	21
Spiders	22
Scorpions	23
Acarina	26
Ticks	38
Mites	41
Silverfish	45
Collembola.	46
Cockroaches	55
Termites	63
Earwigs	64
Psocids	65
Lice (Anoplura)	66
Lice (Mallophaga)	92
Bugs.	94
Lepidoptera	96
Beetles	98
Hymenoptera	100
Flies	120
Mosquitoes.	134
Fleas	167
Snakes	175
Birds	179
Rodents.	181
Lagomorphs	185
Bats.	186
Selected References	187



PUBLIC HEALTH SIGNIFICANCE OF GROUPS INCLUDED IN THE KEY

<i>COMMON NAME</i>	<i>PUBLIC HEALTH SIGNIFICANCE</i>
Ant	bite, sting; infest stored food; damage wood.
Bat	associated with rabies, histoplasmosis and many other diseases.
Bed Bug	cause dermatitis; not known to transmit disease.
Bee, Hornet, etc.	bite and sting; infest stored food; damage wood.
Beetle	infest stored food; infest human intestine; cause dermatitis.
Bird	associated with histoplasmosis, ornithosis and many other diseases.
Book Louse, Psocid	infest stored food.
Caterpillar	sting; infest intestinal tract.
Centipede	venomous bite; infest nasal, intestinal, and urinary tracts.
Chewing Louse	infest domestic birds and mammals.
Cockroach	transmit enteric diseases.
Collembola	infest stored food; used as indicator organisms for pesticide studies.
Copepod	involved in transmission of broad fish tapeworm and guinea worm.
Daddy Long-leg Spider	infest houses; harmless.
Earwig	household pests.
Flea	cause dermatitis; transmit plague, murine typhus, tapeworms.
Fly	some bite; larvae infest human flesh; transmit typhoid, paratyphoid, cholera, bacillary dysentery, infantile diarrhea, amebic dysentery, giardiasis, helminths, trachoma, conjunctivitis, yaws, anthrax, tularemia, African sleeping sickness, leishmaniasis, onchocerciasis, loiasis, bartonellosis, sandfly fever.
Ked or Louse Fly	occasionally bite man.
Kissing Bug	transmit Chagas disease.
Lagomorph	transmit tularemia and many other diseases.
Lobster, Crab, etc.	involved in transmission of oriental lung fluke.
Millipede	exude vesicating venom; infest digestive and urinary tract; intermediate host of tapeworms.
Mite	cause dermatitis; infest human intestine; transmit scrub typhus, rickettsialpox, epidemic hemorrhagic fever.
Mosquito	transmit malaria, encephalitis, yellow fever, dengue, filariasis.
Moth or Butterfly	infest stored food; infest human intestine; some have stinging hairs.
Pseudoscorpion	infest houses; harmless.
Rodent	transmit leptospirosis, lymphocytic choriomeningitis, etc.
Scorpion	sting.
Sea Spider	appearance causes fear; harmless.
Silverfish, Firebrat	infest stored food; transmit enteric diseases.
Snake	venomous bite; secondary infection of bites.
Sowbug, Pillbug	household pests; harmless.
Spider	venomous bite.
Sucking Louse	cause dermatitis; transmit epidemic typhus, trench fever, relapsing fever.
Sun Spider	non-venomous bite.
Termite	destroy wood; housing deterioration.
Thrips	bite man occasionally.
Tick	cause dermatitis, tick paralysis; transmit spotted fever, relapsing fever, tularemia, Colorado tick fever, Russian spring-summer encephalitis.
Whip Scorpion	appearance causes fear; harmless.

INTRODUCTION

Public health biologists are often responsible for teaching animal identification to personnel (sanitarians, engineers, physicians, veterinarians, etc.) without special training in taxonomy. One of the most successful devices for such training has been the pictorial key. The first U.S. Public Health Service pictorial key was devised by Stanley B. Freeborn and Eugene J. Gerberg (1943) to guide personnel in the identification of anopheline mosquito larvae during our national malaria control program.

After the Communicable Disease Center was founded (1946) additional keys were developed. At present the Center utilizes more than 75 such keys in its regular training program. These are the major items incorporated into this booklet. Apropos morphological diagrams are also included.

Precise identification of disease vectors is essential to their efficient control. In using the following keys it should be remembered that only a few of them include all species in a group, and that determinations made using them are only tentative.

The pictorial keys are typical of identification keys found in reference works and scientific papers except that they are arranged as diagrams and are illustrated. After making the first choice offered at the top of each page, follow the black lines or indicated numbers to secondary choices until the correct identification has been made. Note that, in some cases, the identification can be made in the first choice.

Note: The differing formats and typography in this publication were deliberately selected to:

- (1) Provide a broad spectrum of taxonomic experience;
- (2) Avoid the stultifying effect of monotonous repetition.

ARTHROPODS OF PUBLIC HEALTH IMPORTANCE: KEY TO COMMON CLASSES AND ORDERS

Harold George Scott and Chester J. Stojanovich

1. Three or 4 pairs of walking legs (Fig. 1 A & B).....2

Five or more pairs of walking legs (Fig. 1 C & D).....33




Fig. 1 A




Fig. 1 B




Fig. 1 C




Fig. 1 D

2. Three pairs of walking legs (Fig. 2 A).....3

Four pairs of walking legs (Fig. 2 B).....25




Fig. 2 A




Fig. 2 B

3. Wings present, well developed (Fig. 3 A).....4

Wings absent or rudimentary (Fig. 3 B & C).....13




Fig. 3 A




Fig. 3 B




Fig. 3 C

4. With one pair of membranous wings (Fig. 4 A). ORDER DIPTERA.....5

With two pairs of wings (Fig. 4 B & C).....6




Fig. 4 A




Fig. 4 B




Fig. 4 C

5. Wings with scales (Fig. 5 A). FAMILY CULICIDAE.....MOSQUITO

Wings without scales (Fig. 5 B). DIPTERA OTHER THAN MOSQUITOES.....FLY




Fig. 5 A




Fig. 5 B

6. Mouthparts adapted for sucking, with elongate proboscis (Fig. 6 A).....7

Mouthparts adapted for chewing, without elongate proboscis (Fig. 6 B).....9




Fig. 6 A




Fig. 6 B

7. Wings densely covered with scales; proboscis coiled (Fig. 7 A). ORDER LEPIDOPTERA.....MOTH OR BUTTERFLY

Wings not covered with scales; proboscis not coiled (Fig. 7 B).....8




Fig. 7 A




Fig. 7 B

8. Wing with fringe of long hair (Fig. 8 A). ORDER THYSANOPTERA.....THRIPS

Wing without long hair (Fig. 8 B). ORDER HEMIPTERA.....KISSING BUG




Fig. 8 A




Fig. 8 B

9. Both pair of wings membranous and similar in structure (Fig. 9 A).....10

Front pair of wings shell-like or leathery, serving as covers for the second pair (Fig. 9 B).....11




Fig. 9 A




Fig. 9 B

10. Both pairs of wings similar in size (Fig. 10 A). ORDER ISOPTERA.....TERMITE

Hind wing much smaller than front wing (Fig. 10 B). ORDER HYMENOPTERA.....BEE, HORNET, WASP, YELLOW JACKET, OR ANT




Fig. 10 A




Fig. 10 B

11. Front wings horny or leathery, without distinct veins (Fig. 11 A).....12

Front wings leathery or paper-like, with distinct veins (Fig. 11 B). ORDER ORTHOPTERA.....COCKROACH




Fig. 11 A




Fig. 11 B

12. Abdomen with prominent cerci; wings shorter than abdomen (Fig. 12 A). ORDER DERMAPTERA.....EARWIG

Abdomen without prominent cerci; wings covering abdomen (Fig. 12 B). ORDER COLEOPTERA.....BEETLE




Fig. 12 A




Fig. 12 B

13. Mouthparts with jaws for chewing (Fig. 13 A).....14
 Mouthparts with a long beak or stylets for sucking up food (Fig. 13 B).....21




Fig. 13 A




Fig. 13 B

14. With three long terminal tails (Fig. 14 A). ORDER THYSANURA.....SILVERFISH AND FIREBRAT
 Without three long terminal tails (Fig. 14 B).....15




Fig. 14 A




Fig. 14 B

15. Abdomen with prominent pair of cerci (Fig. 15 A). ORDER DERMAPTERA.....EARWIG
 Abdomen without prominent pair of cerci (Fig. 15 B).....16




Fig. 15 A



Fig. 15 B

16. With narrow waist (Fig. 16 A). ORDER HYMENOPTERA.....ANT
 Without narrow waist (Fig. 16 B).....17




Fig. 16 A




Fig. 16 B

17. Antenna with fewer than 8 segments (Fig. 17 A).....18

Antenna with more than 8 segments (Fig. 17 B).....19



Fig. 17 A




Fig. 17 B

18. Abdomen with 6 or fewer segments (Fig. 18 A). ORDER COLLEMBOLA.....SPRINGTAIL

Abdomen with more than 6 segments (Fig. 18 B). ORDER MALLOPHAGA.....CHEWING LOUSE




Fig. 18 A




Fig. 18 B

19. Tarsus with 4-5 segments (Fig. 19 A).....20

Tarsus with 1-3 segments (fig. 19 B). ORDER PSOCOPTERA.....BOOK LOUSE OR PSOCID




Fig. 19 A




Fig. 19 B

20. Pronotum narrower than head, never covering head (Fig. 20 A). ORDER ISOPTERA.....TERMITE

Pronotum broader than head, often covering head (Fig. 20 B). ORDER ORTHOPTERA.....COCKROACH




Fig. 20 A




Fig. 20 B

21. Flattened laterally (Fig. 21 A). ORDER SIPHONATERA.....FLEA

Flattened dorso-ventrally (Fig. 21 B).....22




Fig. 21 A




Fig. 21 B

22. Foot terminating in protrusible bladder (Fig. 22 A). ORDER THYSANOPTERA.....THRIPS

Foot not terminating in protrusible bladder (Fig. 22 B).....23




Fig. 22 A




Fig. 22 B

23. Beak jointed (Fig. 23 A). ORDER HEMIPTERA.....BEDBUG

Beak not jointed (Fig. 23 B).....24




Fig. 23 A




Fig. 23 B

24. Mouthparts retracted into head (Fig. 24 A). ORDER ANOPLURA.....SUCKING LOUSE

Mouthparts not retracted into head (Fig. 24 B). ORDER DIPTERA.....KED OR LOUSE FLY




Fig. 24 A




Fig. 24 B

25. Abdomen well-developed (Fig. 25 A). CLASS ARACHNIDA.....26

Abdomen peg-like (Fig. 25 B). CLASS PYCNOGONIDA.....SEA SPIDER




Fig. 25 A




Fig. 25 B

26. Abdomen distinctly segmented (Fig. 26 A).....27

Abdomen not distinctly segmented (Fig. 26 B).....31




Fig. 26 A




Fig. 26 B

27. Abdomen lengthened to form a long tail (Fig. 27 A).....28

Abdomen not lengthened to form a long tail (Fig. 27 B).....29




Fig. 27 A




Fig. 27 B

28. Tail with stinger (Fig. 28 A). ORDER SCORPIONIDA.....SCORPION

Tail without stinger (Fig. 28 B). ORDER PEDIPALPIDA.....WHIP SCORPION




Fig. 28 A




Fig. 28 B

29. With large pincer-like claws (Fig. 29 A). ORDER PSEUDOSCORPIONIDA.....PSEUDOSCORPION
 Without large pincer-like claws (Fig. 29 B).....30




Fig. 29 A




Fig. 29 B

30. Legs not longer than body (Fig. 30 A). ORDER SOLPUGIDA.....SUN SPIDER
 Legs much longer than body (Fig. 30 B). ORDER PHALANGIDA.....DADDY LONG-LEG SPIDER




Fig. 30 A




Fig. 30 B

31. Abdomen constricted to form a narrow waist (Fig. 31 A). ORDER ARANEIDA.....SPIDER
 Abdomen not constricted (Fig. 31 B).....32




Fig. 31 A




Fig. 31 B

32. Body with long hair; Haller's organ absent (Fig. 32 A). ORDER ACARINA.....MITE
 Body without hair or short hair; Haller's organ present (Fig. 32 B). ORDER ACARINA.....TICK







Fig. 32 A





Haller's organ

Fig. 32 B




33. Five to 7 pairs of walking legs (Fig. 33 A). CLASS CRUSTACEA.....34
More than 14 pairs of walking legs (Fig. 33 B).....36





34. Abdomen without appendages (Fig. 34 A). ORDER COPEPODA.....COPEPOD
Abdomen with appendages (Fig. 34 B).....35




35. Thorax covered with a fused plate; eyes, when present, on movable stalks (Fig. 35 A & B).....
ORDER DECAPODA.....LOBSTER, CRAB, CRAYFISH, SHRIMP, ETC.
Thorax not covered with a fused plate; eyes, when present, not on movable stalks (Fig. 35 C & D)....
ORDER ISOPODA.....SOWBUG, PILLBUG




36. One pair of legs per body segment (Fig. 36 A). CLASS CHILOPODA.....CENTIPEDE
Two pairs of legs per body segment (Fig. 36 B). CLASS DIPLOPODA.....MILLIPEDE




HOUSEHOLD AND STORED-FOOD PESTS: PICTORIAL KEY TO COMMON LARVAE
 Chester J. Stojanovich & Harold George Scott




HOUSEHOLD AND STORED-FOOD PESTS: KEY TO COMMON ADULTS
Harold George Scott & Chester J. Stojanovich




ECTOPARASITES OF THE DOG: PICTORIAL KEY TO COMMON SPECIES
Harold George Scott & Chester J. Stojanovich




REPRESENTATIVE ECTOPARASITES OF THE DOG
Chester J. Stojanovich




Trichodectes canis DOG BITING LOUSE




Linognathus setosus DOG SUCKING LOUSE




Ctenocephalides felis CAT FLEA



Sarcoptes scabiei canis SARCOPTIC MANGE MITE




Otobius megnini SPINOSE EAR TICK



Rhipicephalus sanguineus BROWN DOG TICK

HUMAN ECTOPARASITES: KEY TO COMMON GROUPS
Chester J. Stojanovich and Harold George Scott



CRUSTACEA: KEY TO SOME MAJOR ORDERS
Chester J. Stojanovich and Harold George Scott

- | | |
|--|---|
| 1. With abdominal appendages (Fig. 1 A)..... | 2 |
| Without abdominal appendages (Fig. 1 B)..... | 7 |




Fig. 1 A




Fig. 1 B

- | | |
|-------------------------------------|---|
| 2. Carapace present (Fig. 2 A)..... | 3 |
| Carapace absent (Fig. 2 B)..... | 6 |




Fig. 2 A




Fig. 2 B

- | | |
|--|------------------|
| 3. With dorsal shield (Fig. 3 A). SHIELD SHRIMP..... | Order NOTOSTRACA |
| Without dorsal shield (Fig. 3 B)..... | 4 |




Fig. 3 A






Fig. 3 B


4. With bivalve shell (Fig. 4 A). SHELL SHRIMP.....Order CONCHOSTRACA
 Without bivalve shell (Fig. 4 B).....5



5. First pleopod rudimentary (Fig. 5 A). OPOSSUM SHRIMP.....Order MYSIDACEA
 First pleopod well-developed (Fig. 5 B, C & D). SHRIMP, CRAYFISH, LOBSTERS, CRABS.....
Order DECAPODA



6. Body laterally compressed (Fig. 6 A). SAND FLEAS, ETC.....Order AMPHIPODA
 Body dorso-ventrally compressed (Fig. 6 B). SOWBUGS, PILLBUGS, ETC.....Order ISOPODA



7. Body not completely enclosed in a bivalve shell (Fig. 7 A). 8

Body completely enclosed in a bivalve shell (Fig. 7 B). OSTRACODS Order PODOCOPA




Fig. 7 A





Fig. 7 B



interior view

8. Body segmented (Fig. 8 A). 9

Body not segmented (Fig. 8 B). WATER FLEAS Order CLADOCERA




Fig. 8 A




Fig. 8 B

9. Eyes stalked (Fig. 9 A). FAIRY SHRIMP Order ANOSTRACA

Eyes not stalked (Fig. 9 B). COPEPODS Order EUCOPEPODA




Fig. 9 A






Fig. 9 B

CENTIPEDES: KEY TO SOME IMPORTANT UNITED STATES SPECIES

Harold George Scott




Scutigera cleopatra



Scolopendra heros

MILLIPEDES: KEY TO SOME IMPORTANT UNITED STATES SPECIES
Harold George Scott, Ph.D.

1. 20-21 body segments 2
More than 29 body segments 3
2. Legs with basal spines *Pleurolomia butleri* (= *Fontaria virginiensis*)
Legs without basal spines *Pseudopolydesmus serratus*




Narceus americanus

3. Body segment 3 with legs *Narceus americanus* (= *Spirobolus marginatus*)
Body segment 3 without legs *Brachyiulus pusillus* (= *Julus virgatus*)

Brachyiulus pusillus



ARACHNIDA: KEY TO COMMON ORDERS OF PUBLIC HEALTH IMPORTANCE
 Harold George Scott & Chester J. Stojanovich



SPIDERS: KEY TO SOME IMPORTANT UNITED STATES SPECIES

Harold George Scott & Chester J. Stojanovich

1. Fangs projecting horizontally (Fig. 1 A). (abdomen without tergites; tarsus with claw tufts and 2 claws)
..... Dugesiella hentzi and others, TARANTULAS
- Fangs projecting vertically (Fig. 1 B)..... 2




Fig. 1 A





Fig. 1 B

2. Six eyes in 3 pairs; fiddle-shaped marking on cephalothorax (Fig. 2 A).
Loxosceles reclusa..... BROWN RECLUSE SPIDERS

Eight eyes (shiny black with red spots; usually with red hourglass on underside of abdomen) (Fig. 2 B).
Latrodectus mactans..... BLACK WIDOW SPIDER




Fig. 2 A






Fig. 2 B


SCORPION DIAGRAM: DORSAL VIEW OF CENTRUROIDES VITTATUS,
Chester J. Stojanovich



COLLEMBOLA: PICTORIAL KEY TO NEARCTIC GENERA
 Harold George Scott, Ph.D.
 SUBFAMILY NEANURINAE




COLLEMBOLA: PICTORIAL KEY TO NEARCTIC GENERA
Harold George Scott, Ph.D.
SUBFAMILY ISOTOMINAE - Part A



COLLEMBOLA: PICTORIAL KEY TO NEARCTIC GENERA

SUBFAMILY ISOTOMINAE - Part B

continued from part A




COLLEMBOLA: PICTORIAL KEY TO NEARCTIC GENERA
Harold George Scott, Ph.D.


SUBFAMILIES TOMOCERINAE AND ENTOMOBRYINAE

TOMOCERINAE

ENTOMOBRYINAE




COLLEMBOLA: PICTORIAL KEY TO NEARCTIC GENERA
Harold George Scott, Ph.D.
FAMILY SMINTHURIDAE




COCKROACHES: PICTORIAL KEY TO SOME COMMON SPECIES

Harry D. Pratt



U.S. DEPARTMENT OF
HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
Communicable Disease Center
Training Branch
Atlanta, Georgia - 1953

COCKROACHES: KEY TO EGG CASES OF COMMON DOMESTIC SPECIES
Harold George Scott, Ph.D. and Margery R. Borom



COCKROACHES: KEY TO SOME COMMON SPECIES FOUND IN THE UNITED STATES
Harry D. Pratt & Chester J. Stojanovich

1. Middle and hind femora both with numerous strong spines along the ventral margin (Fig. 1 A)...2
 Middle and hind femora without strong spines along the ventral margin (Fig. 1 B).....12




Fig. 1 A




Fig. 1 B

2. Comparatively large species 18 mm. or longer; subgenital plate of female divided longitudinally, valvular (Fig. 2 A); male styli similar, slender, elongate and straight (Fig. 2 B).....3

Species usually less than 18 mm. long; or, if longer, anterior-ventral margin of front femur with several large stout spines on basal portion, followed by a row of smaller spines (Fig. 2 C); female subgenital plate simple, not divided (Fig. 2 D); male styli variable, frequently modified, asymmetrical, or unequal in size (Fig. 2 E).....8




Fig. 2 A




Fig. 2 D




Fig. 2 B




Fig. 2 C




Fig. 2 E

3. Front wing in both sexes extending beyond tip of abdomen (Fig. 3 A).....4
Front wing in both sexes not reaching tip of abdomen (Fig. 3 B).....7




Fig. 3 A




Fig. 3 B

4. Uniformly dark blackish-brown, shining species (Fig. 4 A).....
.....(Periplaneta fuliginosa) SMOKY BROWN COCKROACH
Species with some yellowish markings on pronotum or front wing or both (Fig. 4 B).....5




Fig. 4 A




Fig. 4 B

5. Front wing with yellowish stripe; pronotum with yellowish and darker areas very contrastingly marked (Fig. 5 A).....(Periplaneta australasiae) AUSTRALIAN COCKROACH
Front wing entirely brownish; pronotum with yellowish and darker areas less contrastingly marked (Fig. 5 B).....6




Fig. 5 A





Fig. 5 B


6. Styli very long and slender, longer than space between their bases (Fig. 6 A); cercus long and slender particularly in the male; male supra-anal plate deeply notched (Fig. 6 B).....
.....(Periplaneta americana) AMERICAN COCKROACH

Styli shorter, not as long as space between their bases (Fig. 6 C); cercus stouter and more evenly spindle-shaped; male supra-anal plate truncate or feebly notched (Fig. 6 D).....
.....(Periplaneta brunnea) BROWN COCKROACH



7. Blackish species, 15-27 mm. long; male front wings covering two-thirds of abdomen (Fig. 7 A); female front wings widely separated pads (Fig. 7 B); first segment of hind tarsus longer than segments 2-5 combined, pulvilli of second and third segments small (Fig. 7 C).....
.....(Blatta orientalis) ORIENTAL COCKROACH

Mahogany brownish species, 30-40 mm. long; front wings reduced to short pads, not widely separated (Fig. 7 D); first segment of hind tarsus shorter than segments 2-5 combined, pulvilli of second and third segments large (Fig. 7 E)....(Eurycotis floridana) LARGE FLORIDA COCKROACH



8. Pronotum with two conspicuous longitudinal dark bars on a pale background (Fig. 8 A).....9

Pronotum variously marked, but without two conspicuous dark longitudinal bars (Fig. 8 B)....10




Fig. 8 A




Fig. 8 B

9. Face pale (Fig. 9 A); male subgenital plate asymmetrical, styli very unequal, short and rounded (Fig. 9 B).....(Blattella germanica) GERMAN COCKROACH

Face dark; male subgenital plate almost symmetrical, styli somewhat elongate and subequal in size (Fig. 9 C).....(Blattella vaga) FIELD COCKROACH




Fig. 9 A

Fig. 9 B




Fig. 9 C

10. Pronotum with a broad dark central stripe; front wings of both sexes appearing to have two transverse brownish bars, some pale specimens showing bars poorly (Fig. 10 A). Width of pronotum usually not exceeding 4.5 mm.....(Supella supellectilium) BROWN-BANDED COCKROACH

Pronotum and front wings otherwise, or, if pronotum is so marked, its width exceeding 4.5 mm.11




Fig. 10 A




Fig. 10 B

11. Larger species 9-25 mm. or more in length; front wing without small dark spots in winged specimens (Fig. 11 A); claws equal (Fig. 11 B); ventral anterior margin of front femur with 3 long apical spines (Fig. 11 C).....(Parcoblatta species) WOOD COCKROACHES

Small species, 8-9 mm. long; front wing with small dark spots (Fig. 11 D); claws unequal (Fig. 11 E); ventral anterior margin of front femur with 2 long apical spines (Fig. 11 F)...
.....(Ectobius pallidus) SPOTTED MEDITERRANEAN COCKROACH




Fig. 11 A

Fig. 11 B




Fig. 11 C




Fig. 11 D




Fig. 11 E




Fig. 11 F

12. Top of eyes close together (Fig. 12 A); general color a nearly uniform greenish; posterior margin of pronotum somewhat angularly produced (Fig. 12 B) (Panchlora nivea) CUBAN COCKROACH

Top of eyes sometimes distant (Fig. 12 C); general color various shades of brown and gray; pronotum usually not angularly produced posteriorly (Fig. 12 D).....,.....13




Fig. 12 A




Fig. 12 B




Fig. 12 C




Fig. 12 D

13. Medium sized species, 30 mm. or less in length, including folded wings (Fig. 14 A & B).....14

Large species 40 mm. or more in length, including folded wings (Fig. 15 A & C).....15

14. Pronotum uniformly blackish except a narrow yellowish band along anterior and lateral margins (Fig. 14 A).....(Pycnoscelus surinamensis) SURINAM COCKROACH

Pronotum pale with a narrow dark longitudinal submarginal band on each side and irregular brownish blotches on disc (Fig. 14 B).....(Nauphoeta cinerea) CINEREOUS COCKROACH




Fig. 14 A




Fig. 14 B

15. Disc or pronotum with shield-like design, sometimes skull-like design (Fig. 15 A); front femur with one or more stout spurs on underside (Fig. 15 B).....
.....(Blaberus giganteus; Blaberus craniifer) GIANT COCKROACH

Disc of pronotum with shield-like design darkened in outline only, not solid black (Fig. 15 C); front femur with a line of stiff hairs on anterior-ventral margin (Fig. 15 D).....
.....(Leucophaea maderae) MADEIRA COCKROACH




Fig. 15 A




Fig. 15 B




Fig. 15 C




Fig. 15 D

TERMITES: KEY TO SOME COMMON NORTH AMERICAN SPECIES
Harold George Scott



Fig. A - Winged Adult



Fig. B - Soldier




Fig. C - Worker

Key to Winged Adults

1. Radius without branches; fontanel (fig. E) usually present 2
Radius (fig. D) with branches; fontanel absent 4
2. Tibia (fig. F) slightly to plainly blackish 3
Tibia entirely pale; Ontario to Guatemala, west to Utah and Arizona
(*Reticulitermes flavipes*) EASTERN SUBTERRANEAN TERMITE
3. Tibia slightly darkened; length 9 mm.; British Columbia to Baja California,
east to Idaho and Sonora
(*Reticulitermes hesperus*) WESTERN SUBTERRANEAN TERMITE
Tibia generally darkened; length 9.5-10 mm.; Oregon and Montana to western
Mexico, Missouri, and Texas
(*Reticulitermes tibialis*) ARID SUBTERRANEAN TERMITE
4. Ocelli (fig. E) present 5
Ocelli absent; western Canada to Baja California
(*Zootermopsis angusticollis*) WESTERN ROTTEN-WOOD TERMITE
5. Body yellow to light brown 6
Body blackish; California to Baja California, east to Arizona and Utah
(*Kalotermes minor*) WESTERN DRY-WOOD TERMITE
6. Transverse rows of long hairs on tergites; South Carolina to Florida,
west to eastern Texas (*Kalotermes snyderi*) EASTERN DRY-WOOD TERMITE
No transverse rows of hairs on tergites; Arizona and California
(*Procryptotermes hubbardi*) ARID DRY-WOOD TERMITE




Fig. D - Wing




Fig. E - Head




Fig. F - Leg

Fig. G - Throat


Key to Soldiers

1. Fontanel (fig. E) present; eyes usually absent 2
Fontanel absent; eyes (fig. E) present 4
2. Gula (fig. G) not twice as broad in front as in middle ARID SUBTERRANEAN TERMITE
Gula twice as broad in front as in middle 3
3. Head twice as long as broad WESTERN SUBTERRANEAN TERMITE
Head less than twice as long as broad EASTERN SUBTERRANEAN TERMITE
4. Antenna (fig. E) with 23-31 segments 5
Antenna with 10-20 segments WESTERN ROTTEN-WOOD TERMITE
5. Third antennal segment as long as next 3 combined EASTERN DRY-WOOD TERMITE
Third antennal segment shorter than next 3 combined WESTERN DRY-WOOD TERMITE
Third antennal segment as long as next 4 combined ARID DRY-WOOD TERMITE

EARWIGS: PICTORIAL KEY TO COMMON DOMESTIC SPECIES

Chester J. Stojanovich and Harold George Scott


tarsus II prolonged beneath III





EUROPEAN EARWIG
Forficula auricularia

wings present

tarsus II not prolonged beneath III




wings absent




SHORE EARWIG
Labidura riparia

legs and antennae banded

legs and antennae not banded




RING-LEGGED EARWIG
Euborellia annulipes



SEASIDE EARWIG
Anisolabis maritima


PSOCIDS: KEY TO SOME SPECIES COMMONLY INFESTING STORED FOOD
 Harold George Scott and Chester J. Stojanovich

- | | |
|---|------------------|
| 1. Two distinct thoracic segments..... | 2 |
| Three distinct thoracic segments (<i>Trogium pulsatorium</i>)..... | DEATH WATCH |
| 2. Without large pronotal bristles | 3 |
| With large pronotal bristles | 4 |
| 3. Eye with 7 facets; head and body brown (<i>Liposcelis bostrychopilus</i>)..... | BANDED PSOCID |
| Eye with 2-4 facets; head brown, body yellow (<i>Liposcelis paetus</i>)..... | WAREHOUSE PSOCID |



Warehouse Psocid


- | | |
|---|--------------|
| 4. Two to 5 large pronotal bristles (<i>Liposcelis entomophilus</i>)..... | GRAIN PSOCID |
| One large pronotal bristle (<i>Liposcelis terricolus</i>) | BOOK LOUSE |



Book Louse

Fig. 140 LICE COMMONLY FOUND ON MAN
Harry D. Pratt

BODY LOUSE
AND
HEAD LOUSE




*PEDICULUS
HUMANUS*


*PHTHIRUS
PUBIS*

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE, Communicable Disease Center, Training Branch, Atlanta, Georgia - Revised Aug. 1953, Oct. 1948

**ANOPLURA: PICTORIAL KEY TO SPECIES ON DOMESTIC RATS
IN SOUTHERN UNITED STATES**
Roy F. Fritz and Harry D. Pratt



ANOPLURA: PICTORIAL KEY TO SOME COMMON GENERA OF SUCKING LICE
Chester J. Stojanovich and Harry D. Pratt



ANOPLURA: KEY TO NORTH AMERICAN SPECIES
Chester J. Stojanovich and Harry D. Pratt

Key to Families of Anoplura

1. Head and thorax more or less thickly covered with setae; in some species the setae are modified into scales (Fig. 1 A). On marine animals.....**FAMILY ECHINOPHTHIRIIDAE**
 Head and thorax with only a few setae (Fig. 1 B).....2

Fig. 1 A





Fig. 1 B



2. Eyes present or with prominent ocular points (Fig. 2 A & B).....3
 Eyes and ocular points absent (Fig. 2 C).....4

Fig. 2 A




Fig. 2 B





Fig. 2 C



3. Abdomen without irregular sclerotized plates on dorsum and venter (Fig. 3 A). On man.....**FAMILY PEDICULIDAE**
 Abdomen with irregular sclerotized plates on dorsum and venter (Fig. 3 B). On hoofed animals.....**FAMILY HAEMATOPINIDAE**

Fig. 3 A





Fig. 3 B



4. Paratergal plates absent (Fig. 4 A). On hoofed animals or carnivores.....
FAMILY LINOGNATHIDAE
 Paratergal plates present (Fig. 4 B). On rodents and lagomorphs...FAMILY HOPLOPLEURIDAE




Fig. 4 A




Fig. 4 B

Key to Genera of Echinophthiriidae

1. Antennae four-segmented; abdomen without scale-like setae (Fig. 1 A).....2
 Antennae five-segmented; abdomen with scale-like setae (Fig. 1 B).....*Antarctophthirus*




Fig. 1 A




Fig. 1 B

2. Legs all essentially the same size (Fig. 2 A).....*Echinophthirius horridus* (von Olfers)
 Anterior legs small; second and third legs stout (Fig. 2 B).....
*Proechinophthirus fluctus* (Ferris)




Fig. 2 A




Fig. 2 B

Key to Species of *Antarctophthirus*

1. Scale-like setae present only on abdomen (Fig. 1 A). *Antarctophthirus callorhini* (Osborn)
 Scale-like setae present on thorax and abdomen (Fig. 1 B) 2




Fig. 1 A




Fig. 1 B

2. Thoracic sternum with a few long setae on posterior border (Fig. 2 A)
 *Antarctophthirus microchir* (Troussart & Neumann)
 Thoracic sternum without long setae on posterior border (Fig. 2 B)
 *Antarctophthirus trichechi* (Bohemann)




Fig. 2 A




Fig. 2 B

Key to Genera of Haematopinidae

1. Sternal plate of thorax present; eyes absent but with prominent ocular points (Fig. 1 A) *Haematopinus*

Sternal plate of thorax absent; eyes present (Fig. 1 B). On peccary..... *Pecaroecus javalii* Babcock & Ewing




Fig. 1 A




Fig. 1 B

Key to Species of *Haematopinus*

1. Thoracic sternal plate wider than long, sternal pits on plate (Fig. 1 A). Hog louse.... *Haematopinus suis* (Linnaeus)

Thoracic sternal plate longer than wide; sternal pits off plate (Fig. 1 B)..... 2




Fig. 2 A




Fig. 2 B

2. Head at least two times as long as wide at ocular points; sternal plate without a median projection (Fig. 2 A & B). On equines. Horse sucking louse..... *Haematopinus asini* (Linnaeus)

Head not two times as long as wide at ocular points; sternal plate with a median projection (Fig. 2 C & D). On cattle..... 3




Fig. 2 A




Fig. 2 B




Fig. 2 C




Fig. 2 D

3. Thoracic sternal plate with median projection blunt and rounded; male genital plate with six setae (Fig. 3 A & B). Short-nosed cattle louse.....
.....*Haematopinus eurysternus* (Nitzsch)

Thoracic sternal plate with median projection more acute and longer; male genital plate with four setae (Fig. 3 C & D). Cattle tail louse.....
.....*Haematopinus quadripertitus* Fahrenholz




Fig. 3 A




Fig. 3 B




Fig. 3 C



Fig. 3 D

Key to Genera of Hoplopleuridae

1. Paratergal plates very small being merely slightly sclerotized points (Fig. 1 A).....
.....*Haemodipsus*

Paratergal plates on at least one abdominal segment usually as long as, or at least half as long as, the sternal plate (Fig. 1 B).....2




Fig. 1 A




Fig. 1 B

2. First and second pair of legs of the same size and form, both being more slender and smaller than the third pair of legs (Fig. 2 A).....3

First pair of legs smallest of the three pairs; the second pair with stouter claws (Fig. 2 B).....4




Fig. 2 A




Fig. 2 B

3. A pair of small sclerotized plates present on venter of abdominal segment 2 (Fig. 3 A); antennae and head without hook-like processes.....Enderleinellus

Sclerotized plates entirely lacking on venter of abdominal segment 2; antennae and head with hook-like processes (Fig. 3 B).....Microphthirus uncinatus (Ferris)




Fig. 3 A




Fig. 3 B

4. Antennae four-segmented (sometimes appearing three-segmented); bladder-like expansions on third leg (Fig. 4 A & B).....Haematopinoides squamosus Osborn

Antennae five-segmented; bladder-like expansions lacking on third leg (Fig. 4 C).....5




Fig. 4 A




Fig. 4 B



Fig. 4 C

5. First sternite of abdominal segment 3 extended laterally to articulate with its corresponding paratergal plate; this sternite bearing two groups of two or three stout setae (Fig. 5 A).....Hoplopleura

First sternite of abdominal segment 3 never articulating with paratergal plate (Fig. 5 B).....6




Fig. 5 A




Fig. 5 B

6. Paratergal plate 2 completely divided longitudinally, one plate on the dorsum and the other on the venter of the abdomen (Fig. 6 A).....Fahrenholzia

Paratergal plate 2 never completely divided to form two distinct plates (Fig. 6 B)....7




Fig. 6 A



Fig. 6 B

1. Sternal plate of thorax usually pointed posteriorly or, if truncate, always associated with a huge enlargement of the first antennal segment (Fig. 7 A & B).....Polyplax

Sternal plate of thorax usually emarginate posteriorly or sometimes quadrate in shape (Fig. 7 C & D).....Neohaematopinus




Fig. 7 A

Fig. 7 B


Fig. 7 C

Fig. 7 D

Key to Species of Enderleinellus


1. Paratergal plates present on abdominal segments 2-5 (Fig. 1 A).....2

Paratergal plates present on abdominal segments 2-6; abdominal sternites and tergites present in both sexes (Fig. 1 B). On Sciurus.....Enderleinellus nitzschi Fahrenholz




SCORPION DIAGRAM: VENTRAL VIEW OF CENTRUROIDES VITTATUS

Chester J. Stojanovich




SCORPIONS: PICTORIAL KEY TO SOME COMMON UNITED STATES SPECIES
 Chester J. Stojanovich and Harold George Scott



ACARINA: ILLUSTRATED KEY TO SOME COMMON ADULT FEMALE MITES AND ADULT TICKS
 Harry D. Pratt and Chester J. Stojanovich


1. Last segment of first leg with a depression known as Haller's organ; most species with a toothed hypostome on capitulum; size usually over 4 mm. (Fig. 1 A). Ticks 21

Last segment of first leg without such a depression known as Haller's organ; hypostome not toothed; most species less than 4 mm. long (Fig. 1 B). Mites..... 2



2. Respiratory system with a spiracle on each side opening latral to the bases of the 3rd or 4th pair of legs, frequently spiracles leading into slender tubes that extend forward laterally to the bases of the 1st or 2nd pairs of legs (Fig. 2 A). Mesostigmatid Mites. 3

Respiratory system without spiracles, or with spiracles opening near bases of the chelicerae (Fig. 2 B)..... 13



3. Anus surrounded by a plate bearing only 3 setae, one on each side and one behind the anal opening; first tarsus bearing caruncle and claws at tip (Fig. 3 A)..... 4

Anus surrounded by a plate bearing more than 3 setae; first tarsus without caruncle and claws (Fig. 3 B)..... Many species of Macrocheles



4. Anal opening more than its length behind anterior margin of anal plate; chelicerae strongly narrowed apically, needle-like, movable chela absent or extremely small (Fig. 4 A). Genus Dermanyssus 5

Anal opening less than its length or about its length, behind anterior margin of anal plate; chelicerae not narrowed apically and needle-like, shear-like, bearing conspicuous shear-like chelae at tip which may or may not bear teeth (Fig. 4 B). 7




Fig. 4 A




Fig. 4 B

5. Dorsal surface of body with a single plate (Fig. 5 A)..... 6

Dorsal surface of body with two plates, a large anterior plate and a small posterior plate (Fig. 5 B). Dermanyssus sanguineus HOUSE MOUSE MITE




Fig. 5 A




Fig. 5 B

6. Peritreme tube somewhat sinuous and extending anteriorly to a point opposite coxa 2 (Fig. 6 A). Dermanyssus gallinae CHICKEN MITE

Peritreme tube short, extending forward for a distance less than half the diameter of coxa 3 (Fig. 6 B). Dermanyssus americanus AMERICAN BIRD MITE




Fig. 6 A




Fig. 6 B

7. Dorsal plate not covering entire dorsal surface of mite; genito-ventral plate typically narrowed posteriorly behind 4th coxae; chelae on chelicerae without teeth or setae (Fig. 7 A). Genus Ornithonyssus 8

Dorsal plate almost covering entire dorsal surface of mite; genito-ventral plate typically expanded posterior to 4th coxae; one or both chelae of chelicerae with teeth and a seta (Fig. 7 B). Family Laelaptidae 10





Fig. 7 A



genito-ventral plate





Fig. 7 B



8. Sternal plate with anterior and middle pairs of sternal setae on the plate, posterior pair usually just off the plate (Fig. 8 A). On Birds... Ornithonyssus sylviarum NORTHERN FOWL MITE

Sternal plate with the usual three pairs of setae on the plate (Fig. 8 B) 9




Fig. 8 A




Fig. 8 B

9. Dorsal plate narrowed posteriorly; setae in middle dorsal row of plate longer than the distance between their bases (Fig. 9 A). Normally on mammals or man. Ornithonyssus bacoti TROPICAL RAT MITE

Dorsal plate broader posteriorly; setae in middle dorsal row of plate much shorter than the distance between their bases (Fig. 9 B). Normally on birds. Ornithonyssus bursa TROPICAL BIRD MITE




Fig. 9 A




Fig. 9 B

10. Genito-ventral plate with many fine setae; anal plate transverse, wider than long (Fig. 10 A). On domestic rats and a wide variety of wild mammals..... Eulaelaps stabularis

Genito-ventral plate with one to four pairs of **setae**; anal plate longer than wide (Fig. 10 B)..... 11




Fig. 10 A




Fig. 10 B

11. Genito-ventral plate with only a single pair of setae (Fig. 11 A). On domestic rats and mice and a wide variety of mammals and birds..... Haemolaelaps glasgowi..... COMMON RODENT MITE

Genito-ventral plate with four pairs of setae (Fig. 11 B). Normally on domestic rats..12




Fig. 11 A




Fig. 11 B

12. Anal plate contiguous with the genito-ventral plate, anterior margin rounded and fitting into a strong concavity in genito-ventral plate; larger species averaging 1-2 mm. long. (Fig. 12 A). Echinolaelaps echidninus..... SPINY RAT MITE

Anal plate somewhat separated from genito-ventral plat, anterior margin almost straight with definite anterior-lateral corners; small species averaging 0.5-1 mm long (Fig. 12 B). Laelaps nuttalli..... DOMESTIC RAT MITE




Fig. 12 A




Fig. 12 B

13. First pair of legs very long, much longer than other three pairs; anterior margin of body with four distinct flattened scales and somewhat flattened scales on other dorsal surfaces of body (Fig. 13 A). Plant feeders which invade buildings but do not bite man. Bryobia praetiosa CLOVER MITE

First pair of legs not markedly longer than the other three pairs of legs; no flattened scales on body (Fig. 13 B) 14




Fig. 13 A




Fig. 13 B

14. Surface of body without fine parallel lines or folds; tarsi without stalked suckers (Fig. 14 A). Adults never true parasites (Cheese or Flour mites) 15

Surface of body with fine parallel lines or folds; tarsi often provided with stalked suckers (Fig. 14 B). Scabies or mange mites parasitic in all stages, chiefly on vertebrates 16




Fig. 14 A




Fig. 14 B

15. Tarsi tapering markedly to tip (Fig. 15 A) Glycyphagus prunorum

Tarsi not tapering markedly to tip (Fig. 15 B). Many cheese and flour mites which are difficult to separate except with very specialized literature and a reference collection. Genus Tyrophagus, Genus Caloglyphus, Etc.




Fig. 15 A




Fig. 15 B

16. Body elongate, somewhat cigar-shaped and prolonged behind; the abdomen somewhat ringed; legs very short, apparently three-segmented; tiny species less than 1 mm. (Fig. 16 A). In hair follicles or sebaceous glands of mammals.....
Demodex folliculorum..... PORE OR FOLLICLE MITE

Body not prolonged behind and cigar-shaped (Fig. 16 B). Occasionally female grain itch somewhat balloon-shaped; larger species not found in hair follicle or sebaceous glands of mammals..... 17




Fig. 16 A




Fig. 16 B

17. A club-shaped or clavate hair between bases of first and second pairs of legs, body divided into cephalothorax and abdomen, the latter often enormously enlarged (Fig. 17 A)
Pyemotes ventricosus formerly Pediculoides ventricosus..... STRAW ITCH MITE

Setae on cephalothorax normal, no club-shaped or clavate hair between bases of first and second pairs of legs; no distinct division into cephalothorax and abdomen (Fig. 17 B)..... 18




Fig. 17 A




Fig. 17 B

18. Legs short and stubby (Fig. 18 A)..... 20

Legs longer and more slender (Fig. 18 B)..... 19




Fig. 18 A




Fig. 18 B

19. Suckers of tarsi with segmented pedicels (Fig. 19 A). Non-burrowing itch mites on mammals in the genus Psoroptes, a common species causing scabs and crusts in the ears of rabbits is the Psoroptes cuniculi RABBIT EAR MITE

Suckers of tarsi without segmented pedicels (Fig. 19 B)..... Dermatophagoides scheremetewskyi




Fig. 19 A



Fig. 19 B

20. Anal opening on the dorsal surface of the body; dorsal surface of the body with only short, sharp setae (Fig. 20 A)..... Notoedres

Anal opening at tip of body or slightly on ventral side; dorsal surface of body with pointed scales and blunt stout spines (Fig. 20 B). Sarcoptes scabiei SCABIES OR MANGE MITE




Fig. 20 A




Fig. 20 B

21. Capitulum at anterior end of body, visible from above and below; scutum or dorsal shield present, short in female, long in male (Fig. 21 A & B). Family Ixodidae. HARD TICKS....22

Capitulum on under side of body, hidden by body when seen from above though palpi may project anteriorly; scutum absent (Fig. 21 C & D). Family Argasidae.....SOFT TICKS....31




Fig. 21 A

Fig. 21 B




Fig. 21 C

Fig. 21 D

FAMILY IXODIDAE - HARD TICKS

22. Ornate ticks, with some white markings on dorsal shield (Fig. 22 A).....23

Inornate ticks, without white markings on dorsal shield (Fig. 22 B).....28




Fig. 2 A




Fig. 2 B

23. Palpi long, much longer than basis capituli; second segment of palpus about twice as long as wide (Fig. 23A). Genus Amblyomma.....24

Palpi short, about as long as basis capituli; second segment of palpus about as long as wide (Fig. 23 B). Genus Dermacentor.....25




Fig. 23A

III --- II --- I --- palpal segments --- III --- II --- I
basis capituli




Fig. 23 B

basis capituli

24. Next to last segment of second, third, and fourth pairs of legs without paired terminal spurs; female with a distinct pale marking near posterior end of dorsal shield (Fig. 24 A). Amblyomma americanum.....LONE STAR TICK

Next to last segment of second, third, and fourth pairs of legs with long, paired terminal spurs; female with more diffuse markings on dorsal shield (Fig. 24 B).....Amblyomma maculatum.....GULF COAST TICK




Fig. 4 A




Fig. 4 B

25. Spiracular plate without dorsal prolongation (Fig. 25 A). Dermacentor albipictus.....WINTER TICK

Spiracular plate with dorsal prolongation (Fig. 25 B).....26




Fig. 25 A




Fig. 25 B

26. Basis capituli with long cornua (Fig. 26 A). Dermacentor occidentalis. PACIFIC COAST TICK

Basis capituli with short cornua (Fig. 26 B).....27




Fig. 26 A




Fig. 26 B

... Goblets of spiracular plate large and less numerous; Rocky Mountain species. (Fig. 27 A) *Dermacentor andersoni* ROCKY MOUNTAIN WOOD TICK

Goblets of spiracular plate very small and numerous; east of the Rocky Mountains and on the Pacific coast. (Fig. 27 B). *Dermacentor variabilis* AMERICAN DOG TICK




Fig. 27 A




Fig. 27 B

28. Sides of basis capituli laterally produced; distinctly angulate; eyes present on sides of scutum (Fig. 28 A & B) 29

Sides of basis capituli not laterally produced; more or less parallel (Fig. 28 C); eyes absent 30




Fig. 28 A




Fig. 28 B




Fig. 28 C

29. Fore coxa deeply cleft; festoons present; easily seen in unengorged specimens; anal groove distinct in unengorged specimens (Fig. 29 A). (principally on dogs or in houses) *Rhipicephalus sanguineus* BROWN DOG TICK

Fore coxa not deeply cleft; festoons absent; anal groove indistinct (Fig. 29 B). (On cattle and deer). *Boophilus annulatus* CATTLE TICK




Fig. 29 A





Fig. 29 B

30. Second segment of palpus laterally produced; anal groove behind anus, not attaining posterior margins of body (Fig. 30 A & B). Haemaphysalis leporispalustris.....RABBIT TICK

Second segment of palpus not laterally produced; anal groove extending as an inverted U from in front of anus to posterior margins of body (Fig. 30 C).....Genus Ixodes



FAMILY ARGASIDAE - SOFT TICKS

31. Margin of body with a definite sutural line separating dorsal and ventral surfaces; dorsal surface with conspicuous "discs" arranged somewhat in radiating lines (Fig. 31 A)
Argas persicus.....FOWL TICK

Margin of body lacking definite sutural line, thick and rounded (Fig. 31 B).....32




Fig. 31 A




Fig. 31 B

32. Hypostome with well-developed teeth (Fig. 32 A); integument not spinose.....
Genus Ornithodoros.....33

Hypostome of adult vestigial or without effective teeth; integument of nymph (stage usually seen) spinose (Fig. 32 B). Usually on cattle and horses.....
Otobius megnini.....SPINOSE EAR TICK



Fig. 32 A




Fig. 32 B

33. Strong dorsal humps absent on all tarsi (Fig. 33 A).....34

Strong dorsal humps present on tarsi of first, second and third legs (Fig. 33 B).....35




Fig. 33




Fig. 33 B

34. Cheeks absent (Fig. 34 A). Ornithodoros hermsi.....HERMS' RELAPSING FEVER TICK

Cheeks present (Fig. 34 B).....Ornithodoros talaje




Fig. 34 A




Fig. 34 B

35. Eyes present on sides of body above second and third coxae (Fig. 35 A); tarsus of fourth leg with a prominent, pointed subterminal spur (Fig. 35 B).....PAJAROELLO TICK
Ornithodoros coriaceus.....

Eyes absent; tarsus of fourth leg without such subterminal spur (Fig. 35 C).....15




Fig. 35 A





Fig. 35 B

Fig. 35 C


36. Mammillae large, relatively few and not crowded; in mid-dorsal region about 10 per linear mm.; hypostome over 1/2 mm. long, Southeastern United States and Mexico north to Kansas and Florida. Ornithodoros turicata.....RELAPSING FEVER TICK

Mammillae small, crowded, and numerous; in mid-dorsal region about 18 per linear mm.; hypostome less than 1/2 mm. long. Pacific coast and Rocky Mountain states.....Ornithodoros parkeri.....PARKER'S RELAPSING FEVER TICK

TICKS: KEY TO GENERA IN UNITED STATES
Harry D. Pratt




TICKS AND MITES: KEY TO SPECIES COMMONLY INFESTING PIGEONS
 Harold George Scott & Chester J. Stojanovich




TICKS: PICTORIAL KEY TO SOME COMMON SPECIES

Harry D. Pratt

capitulum visible from above,
scutum present, family Ixodidae,
HARD TICKS





capitulum not visible from above,
scutum absent, family Argasidae.
SOFT TICKS



sutural line present



sutural line absent

*Argas persicus*
FOWL TICK*Ornithodoros*
RELAPSING FEVER TICK

mouthparts short, about as long
as basis capituli



mouthparts much longer than basis capituli
white spot on tip of scutum of female

mouthparts basis capitulum


*Amblyomma americanum*
LONE STAR TICK

scutum with white markings; basis
capituli with parallel sides

scutum without white markings; basis
capituli produced laterally to form an angle


*Dermacentor variabilis* and *D. andersoni*
AMERICAN DOG TICK AND WOOD TICK*Rhipicephalus sanguineus*
BROWN DOG TICK

MITE DIAGRAM WITH STRUCTURES LABELED
Harry D. Pratt




MITES: PICTORIAL KEY TO SOME COMMON SPECIES OF PUBLIC HEALTH IMPORTANCE
Harold George Scott and Chester J. Stojanovich

with lateral spiracles and peritreme







without lateral spiracles or peritreme



anal opening at rear of anal plate

anal opening at front of anal plate





CHICKEN MITE

Dermanyssus gallinae

genito-ventral plate narrowed posteriorly

genito-ventral plate expanded posteriorly



TROPICAL RAT MITE

Ornithonyssus bacoti

SPINY RAT MITE

Echinolaelaps echidninus


body oval with fine lines

body strongly elongate

body somewhat elongate

abdomen with fine lines

abdomen without lines




ITCH MITE

Sarcoptes scabiei



FOLLICLE MITE

Demodex folliculorum



A CHEESE MITE

Tyrophagus lintneri


U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
 PUBLIC HEALTH SERVICE, Communicable Disease Center, Training Branch, Atlanta, Georgia

- 1963


A FIELD KEY TO SOME SPECIES COMMONLY INFESTING HOUSEHOLDS AND STORED FOOD

Harold George Scott


1. With club-like hair between bases of legs I and II 5
Without club-like hair between bases of legs I and II 2
2. Claws, if present, not on stalks (*Glycyphagus domesticus*, formerly
Glyciphagus prunorum) SUGAR MITE
Claws on stalks 3
3. Internal apical hair (on joint between femur I and tibia I) less
than three times as long as external apical hair 4
Internal apical hair (on joint between femur I and tibia I) more
than three times as long as external apical hair (*Acarus farinae*,
formerly *Tyroglyphus farinae*) HAM MITE

*Acarus farinae*


4. Tarsus with one stout dorsal and five small ventral terminal spines
(*Acarus siro*, formerly *Tyroglyphus siro*) GRAIN MITE
Tarsus with only three small ventral spines (*Tyrophagus castellani*,
formerly *Tyroglyphus longior*) CHEESE MITE

*Tyrophagus castellani*


5. Tarsus IV of female ending in claws and a fleshy protuberance; leg
IV of male smoothly curved inwards (*Pyemotes ventricosus*, formerly
Pediculoides ventricosus) STRAW ITCH MITE
Tarsus IV of female ending in two long hairs of unequal length; leg
IV of male sharply bent (*Tarsonemus floriculus*) FLORICOLUS GRAIN MITE

*Pyemotes ventricosus*


**MITES: PICTORIAL KEY TO ADULT FEMALES
COMMONLY FOUND ON DOMESTIC RATS IN SOUTHERN UNITED STATES**
Harry D. Pratt and Chester J. Stojanovich




SILVERFISH. PICTORIAL KEY TO DOMESTIC SPECIES
Chester J. Stojanovich and Harold George Scott




COLLEMBOLA: PICTORIAL KEY TO COMMON DOMESTIC SPECIES
 Harold George Scott, and Chester J. Stojanovich



COLLEMBOLA DIAGRAMS
Harold George Scott




SUBORDER ARTHROPLEONA




SUBORDER SYMPHYPLEONA

COLLEMBOLA: PICTORIAL KEY TO WORLD SUBFAMILIES
Harold George Scott, Ph.D.



COLLEMBOLA: PICTORIAL KEY TO NEARCTIC GENERA
Harold George Scott, Ph.D.

SUBFAMILIES PODURINAE, HYPOGASTRURINAE, AND ONYCHIURINAE



2. Paired ventral plates of abdominal segment 2 completely detached from its corresponding paratergal plate; each ventral plate bearing a single seta (Fig. 2 A). On Sciurus.....3

Paired ventral plates of abdominal segment 2 each extending laterally to unite with its corresponding paratergal plate; ventral plates without setae (Fig. 2 B).....5




Fig. 2 A

ventral plate

Fig. 2 B

3. Spermatheca present; arms of basal plate apically bilobed (Fig. 3 A & B).....4

Spermatheca absent; arms of basal plate not apically bilobed (Fig. 3 C).....Enderleinellus kelloggi Ferris




Fig. 3 A

Fig. 3 B

Fig. 3 C

1. Spermatheca a straight slightly tapering tube; arms of basal plate apically bilobed but not expanded (Fig. 4 A & B) Enderleinellus longiceps (Kellogg & Ferris)

Spermatheca bent and with its ends expanded; arms of basal plate apically expanded and strongly bilobed (Fig. 4 C) Enderleinellus arizonensis Werneck




Fig. 4 A

Fig. 4 B

Fig. 4 C

5. Paratergal plate 5 and lateral margin of abdominal segment 6 without a pair of long setae (Fig. 5 A) 6

Paratergal plates or lateral margins of abdominal segments 4-8 with a pair of long setae (Fig. 5 B). On Marmota Enderleinellus marmotae Ferris




Fig. 5 A



Fig. 5 B

6. Female with 2-4 long setae on dorsum of abdominal segment 4 reaching to apex of body (Fig. 5 A). On Citellus and Cynomys Enderleinellus osborni (Kellogg & Ferris)



Female without such setae. On Citellus Enderleinellus suturalis (Osborn)

Key to Species of *Fahrenholzia*

1. Paratergal plates present only on abdominal segments 2 to 4 (Fig. 1 A) 2
 Paratergal plates present on at least abdominal segments 2-6 (Fig. 1 B) 6



2. Dorsal surface of abdomen with a narrow, sclerotized, median, longitudinal plate between paratergal plates 2 (Fig. 2 A). On Liomys 3
 Dorsal surface of abdomen without such a plate (Fig. 2 B). On Perognathus and Dipodomys 5



3. Thoracic sternal plate concave on anterior margin; dorsal lobe of paratergal plate 3 pointed apically (Fig. 3 A & B) *Fahrenholzia texana* Stojanovich & Pratt
 Thoracic sternal plate convex on anterior margin; dorsal lobe of paratergal plate 3 apically truncate (Fig. 3 C & D) 4







Fig. 3 A

Fig. 3 B

Fig. 3 C

Fig. 3 D

4. Dorsal lobe of paratergal plate 2 with the smaller seta about as long as the plate (Fig. 4 A) Fahrenholzia ehrlichi Johnson

Dorsal lobe of paratergal plate 2 with the smaller seta minute, much shorter than the plate (Fig. 4 B) Fahrenholzia microcephala Ferris




Fig. 4 A





Fig. 4 B

5. Paratergal plates of abdominal segment 2 with a single pair of setae between dorsal and ventral lobes; male genitalia with parameres greatly expanded; female genital plate present (Fig. 5 A, B, & C) Fahrenholzia pinnata Kellogg & Ferris

Paratergal plates of abdominal segment 2 with 6 to 8 long setae between dorsal and ventral lobes; parameres of male genitalia not expanded; female genital plate absent (Fig. 5 D & E) Fahrenholzia reducta Ferris



6. Paratergal plates present on abdominal segments 2 to 6; paratergal plate 3 bilobed (Fig. 6 A) Fahrenholzia zacatecae Ferris

Paratergal plates present on abdominal segments 2 to 7; paratergal plate 3 not bilobed (Fig. 6 B) Fahrenholzia tribulosa Ferris




Fig. 6 A




Fig. 6 B

Key to Species of Hoplopleura

1. Third abdominal sternal plate with two groups of two stout setae (Fig. 1 A).....2

Third abdominal sternal plate with two groups of three stout setae (Fig. 1 B).....
On Glaucomys.....Hoplopleura trispinosa Kellogg & Ferris




Fig. 1 A




Fig. 1 B

2. Posterior margins of paratergal plates 3-5 with a broad or pointed lobe on each side (Fig. 2 A & B).....3

Posterior margins of paratergal plates 3-5 with four rounded lobes (Fig. 2 C).....
On Oryzomys.....Hoplopleura oryzomydis Pratt & Lane




Fig. 2 A




Fig. 2 B




Fig. 2 C

3. Paratergal plates 4 and 5 with broad lobes on posterior margin (Fig. 3 A).....4

Paratergal plates 4 and 5 with pointed lobes on posterior margin (Fig. 3 B).....7




Fig. 3 A




Fig. 3 B

4. Faratergal plates 4 and 5 with one large and one minute seta on posterior margin (Fig. 4 A).....5

Faratergal plates 4 and 5 with two large setae on posterior margin (Fig. 4 B).....
On field rodents.....Hoplopleura acanthopus (Burmeister)




Fig. 4 A




Fig. 4 B

5. Abdomen with setae in some of the membrane between sternal and paratergal plates (Fig. 5 A). On Rattus.....Hoplopleura oenomydis Ferris

Abdomen without setae in membrane between ends of sternal and paratergal plates (Fig. 5 B)6




Fig. 5 A




Fig. 5 B

6. Thoracic sternal plate pointed posteriorly (Fig. 6 A). On Peromyscus.....
.....*Hoplopleura hesperomydis (Osborn) and *Hoplopleura ferrisi Cook & Beer

Thoracic sternal plate blunt posteriorly (Fig. 6 B). On Onychomys.....
.....Hoplopleura onychomydis Cook & Beer




Fig. 6 A




Fig. 6 B

7. Thoracic sternal plate about as long as broad; first sternal plate on abdominal segment 3 with two stout setae usually set close together on each side (Fig. 7 A).....8

Thoracic sternal plate definitely longer than broad; first sternal plate on abdominal segment 3 with two stout setae more widely spaced on each side (Fig. 7 B).....9




Fig. 7 A




Fig. 7 B

*These species are separated only in the immature stages.

8. Paratergal plate 6 with posterior angles produced into points (Fig. 8 A). On Eutamias Hoplopleura arboricola Kellogg & Ferris
 Paratergal plate 6 without points on posterior angles (Fig. 8 B). On Tamias Hoplopleura erratica (Osborn)




Fig. 8 A




Fig. 8 B

9. Posterior margin of paratergal plate 6 with angles produced to form a deep emargination (Fig. 9 A). On Sciurus Hoplopleura sciuricola Ferris

Posterior margin of paratergal plate 6 with angles not produced to form a deep emargination (Fig. 9 B). On Sigmodon 10




Fig. 9 A




Fig. 9 B

10. Female with paratergal plates 4-6 elongated; male with 11 tergal plates bearing a row of setae (Fig. 10 A & B) Hoplopleura arizonensis Stojanovich & Pratt

Female with paratergal plates 4-6 only slightly elongated; male with only 7 tergal plates bearing a row of setae (Fig. 10 C & D) Hoplopleura hirsuta Ferris




Fig. 10 A



Fig. 10 C




Fig. 10 B




Fig. 10 D

Key to Species of Haemodipsus

1. Thoracic sternal plate almost three times as wide as long (Fig. 1 A). On domestic rabbits (Oryctolagus).....Haemodipsus ventricosus (Denny)

Thoracic sternal plate hexagonal, being almost as long as wide (Fig. 1 B). On wild rabbits and hares (Sylvilagus and Lepus).....Haemodipsus setoni Ewing




Fig. 1 A




Fig. 1 B

Key to Species of Neohaematopinus

1. Thoracic sternal plate concave on posterior margin (Fig. 1 A).....2

Thoracic sternal plate somewhat oval, and convex on posterior margin (Fig. 1 B).....11




Fig. 1 A




Fig. 1 B




Fig. 1 C

2. Paratergal plates 3 to 6 with three spines on posterior margins (Fig. 2 A).....3

Paratergal plates 3 to 6 with two spines on posterior margins (Fig. 2 B).....5




Fig. 2 A




Fig. 2 B

3. Posterior angle of first antennal segment with a stout spine (Fig. 3 A). On Eutamias...
.....Neohaematopinus pacificus (Kellogg & Ferris)

Posterior angle of first antennal segment without a stout spine (Fig. 3 B).....4



Fig. 3 A




Fig. 3 B

4. Abdominal tergal and sternal plates present on each segment in both sexes (Fig. 4 A)....
On Citellus tereticaudus.....Neohaematopinus citellinus Ferris

Abdominal tergal and sternal plates absent in the middle segments of female; male with
only sternal plates absent (Fig. 4 B). On Citellus spilosoma.....
.....Neohaematopinus spilosomae Stojanovich & Pratt




Fig. 4 A




Fig. 4 B

5. First antennal segment prolonged posterio-apically, with stout spine (Fig. 5 A).....6

First antennal segment without such a prolongation (Fig. 5 B).....8




Fig. 5 A




Fig. 5 B

6. Female without sternal and tergal plates on abdominal segments except for the normal
terminal and genital segments (Fig. 6 A). On Sciurus griseicolor.....
.....Neohaematopinus griseicolor Ferris

Female with sternal and tergal plates on all abdominal segments (Fig. 6 B).....7




Fig. 6 A




Fig. 6 B

7. Second antennal segment with short spine-like seta on posterior margin (Fig. 7 A)
On Tamias hudsonicus Neohaematopinus semifasciatus Ferris

Second antennal segment without spine-like seta (Fig. 7 B). On Sciurus niger
..... Neohaematopinus sciurinus Mjøberg




Fig. 7 A




Fig. 7 B

8. Abdominal sternal and tergal plates absent in female; male with only sternal plates ab-
sent (Fig. 8 A). On Neotoma cinerea Neohaematopinus inornatus Ferris

Abdominal sternal and tergal plates present in both sexes (Fig. 9 A) 9




Fig. 8 A

9. A row of setae present on membrane between most of the sternal and tergal plates of ab-
domen (Fig. 9 A) 10

Membrane between the abdominal sternal and tergal plates without a row of setae (Fig. 9
B). On Glaucomys Neohaematopinus sciuropteri (Osborn)




Fig. 9 A




Fig. 9 B

10. First antennal segment with a spine-like seta at the posterio-apical angle (Fig. 10 A)
On Sciurus carolinensis.....Neohaematopinus sciuri Jancke

First antennal segment with a spine-like seta set somewhat away from the margin in the posterio-apical angle (Fig. 10 B). On Neotoma albicula, streator and micropus.....
.....Neohaematopinus neotomae Ferris




Fig. 10 A




Fig. 10 B

11. Thoracic spiracle small, about one-fourth length of second coxa (Fig. 11 A).....
On Citellus and Cynomys.....Neohaematopinus laeviusculus (Grube)

Thoracic spiracle larger, almost one-half length of second coxa (Fig. 11 B).....
On Marmota.....Neohaematopinus marmotae Ferris




Fig. 11 A




Fig. 11 B

Key to Species of Polyplax

1. Sternal plate of thorax rounded or pointed posteriorly (Fig. 1 A).....2

Sternal plate of thorax truncate posteriorly (Fig. 1 B). On Peromyscus and Onychomys...
.....Polyplax auricularis Ferris




Fig. 1 A




Fig. 1 B

2. Paratergal plate 4 with both setae short or subequal (Fig. 2 A).....3

Paratergal plate 4 with dorsal seta longer than ventral seta; usually as long or longer than plate (Fig. 2 B). On house mouse.....Polyplax serrata (Burmeister)



Fig. 2 A




Fig. 2 B

3. Paratergal plates 3-5 with both apical angles produced into points (Fig. 3 A).....4
On microtene mice.....

Paratergal plates 3-5 with only dorsal apical angle produced into a point (Fig. 3 B)....
On Rattus.....Polyplax spinulosa (Burmeister)




Fig. 3 A




Fig. 3 B

4. First abdominal sternal plate strongly arcuate and with its lateral angles somewhat prolonged (Fig. 4 A).....Polyplax borealis Ferris

First abdominal sternal plate not arcuate, its posterior margin almost straight and lateral angles not produced (Fig. 4 B).....Polyplax alaskensis Ewing




Fig. 4 A




Fig. 4 B

Key to Genera of Linognathidae

1. Sternal plate of thorax at least half as wide as long (Fig. 1 A).....Solenopotes
Sternal plate of thorax small and slender or completely lacking (Fig. 1 B).....Linognathus




Fig. 1 A




Fig. 1 B

Key to Species of Linognathus

1. Head about as broad as long; antennae almost as long as head (Fig. 1 A).....2
Head almost twice as long as wide or longer; antennae noticeably shorter than head (Fig. 1 B).....3




Fig. 1 A




Fig. 1 B

2. Thoracic dorsum with four long setae; head slightly longer than broad (Fig. 2 A). On dogs, foxes and ferrets. Dog sucking louse.....Linognathus setosus (von Olfers)
Thoracic dorsum with two long setae; head definitely as broad as long (Fig. 2 B).....Linognathus pedalis (Osborn)




Fig. 2 A




Fig. 2 B

3. Fore head acutely conical and much elongated; female gonopod with a sclerotized hook (Fig. 3 A & B). On cattle. Long-nosed cattle louse.....*Linognathus vituli* (Linnaeus)

Fore head rounded (Fig. 3 C); female gonopod rounded or with a slight tooth (Fig. 5 B & C). On sheep and goats.....4




Fig. 3 A




Fig. 3 B




Fig. 3 C

4. Head greatly expanded behind antennae; female gonopod rounded (Fig. 4 A & B). Goat sucking louse.....*Linognathus africanus* (Kellogg & Paine)

Head not greatly expanded behind antennae (Fig. 4 C).....5




Fig. 4 A




Fig. 4 B




Fig. 4 C

5. Thoracic spiracle large and conspicuous; female gonopod rounded (Fig. 5 A & B). Sheep louse.....*Linognathus ovillus* (Neumann)

Thoracic spiracle not large and conspicuous; female gonopod with a slight tooth (Fig. 5 C & D). Goat sucking louse.....*Linognathus stenopsis* (Burmeister)




Fig. 5 A




Fig. 5 B




Fig. 5 C




Fig. 5 D

Key to Species of Solenopotes

1. Abdominal spiracles strongly protuberant (Fig. 1 A); female genitalia with apical processes strongly constricted near middle (Fig. 1 B); male genitalia as in figure 2 E. On cattle. Little blue cattle louse.....Solenopotes capillatus Enderlein

Abdominal spiracles only slightly protuberant (Fig. 1 C); female genitalia with apical processes not constricted (Fig. 1 D & E); male genitalia as in figures 2 C & D. On deer.....2

Fig. 1 A





Fig. 1 C

Fig. 1 B (capillatus)Fig. 1 D (binipilosus)Fig. 1 E (ferrisi)

2. Neck present, head with distinct posterior-lateral angles (Fig. 2 A); female genitalia as in figure 1 E; male genitalia as in figure 2 C.....Solenopotes ferrisi (Fahrenholz)

Head without distinct posterior-lateral angles (Fig. 2 B); female genitalia as in figure 1 D ; male genitalia as in figure 2 D.....Solenopotes binipilosus (Fahrenholz)

Fig. 2 A







Fig. 2 B

Fig. 2 C
(ferrisi)Fig. 2 D
(binipilosus)Fig. 2 E
(capillatus)

Key to Genera of Pediculidae

1. Abdomen much longer than basal width; without hairy tubercles (Fig. 1 A). Head and body louse.....Pediculus humanus Linnaeus

Abdomen about as long as basal width; with hairy tubercles (Fig. 1 B). Crab louse....
.....Pthirus pubis (Linnaeus)




Fig. 1 A





Fig. 1 B


MALLOPHAGA: PICTORIAL KEY TO SPECIES INFESTING PIGEONS

Harold George Scott and Chester J. Stojanovich


maxillary palps present



maxillary palps absent




forehead with spines




Hohorstiella lata
LARGE PIGEON BODY LOUSE

forehead without spines
femur III with comb




Colpocephalum turbinatum
SMALL PIGEON BODY LOUSE

femur III without comb



Bonomiella columbae
PIGEON VENT LOUSE

head longer than wide




Columbicola columbae
SLENDER PIGEON LOUSE


head wider than long
forehead with spines




Physconelloides zenaidurae
PIGEON HEAD LOUSE



male basal antennal segment small




male basal antennal segment large



Campanulotes bidentatus compar
SMALL PIGEON FEATHER LOUSE


Coloceras damicornis fahrenholzi
LARGE PIGEON FEATHER LOUSE

MALLOPHAGA: PICTORIAL KEY TO SOME COMMON SPECIES ON CHICKENS
Chester J. Stojanovich and Harry D. Pratt




BUGS: PICTORIAL KEY TO SOME SPECIES THAT MAY BITE MAN
Harry D. Pratt and Chester J. Stojanovich

wings usually well-developed; body elongate-oval



ASSASSIN AND KISSING BUGS-FAMILY REDUVIIDAE


wings reduced; body broadly-oval



BED BUGS-FAMILY CIMICIDAE

thorax with cog wheel crest


thorax without crest



WHEEL BUG
Arilus cristatus

fore-wing with 2 yellow spots


fore-wing dark in U. S. species



CORSAIR
Rasabus biguttatus

antenna inserted midway between eye and tip of head; beak slender, straight


antenna inserted near eye; beak stout, curved




KISSING BUG
Triatomina spp.

pronotum constricted behind middle

pronotum constricted before middle




BLACK BUG
Melanolestes picipes




MASKED HUNTER
Reduvius personatus

middle coxae nearly touching
beak reaching 2nd coxa



POULTRY BUG
Haematosiphon inodus

middle coxae widely separated
beak not reaching 2nd coxa



3rd and 4th antennal segments equal




4th antennal segment shorter than 3rd




BARN SWALLOW BUG
Occlusus picarius

fringe hairs on pronotum longer than, or equal to, width of eye




BAT BUGS
Cimex adjunctus E. N. AM.
Cimex pilosellus W. N. AM.

fringe hairs on pronotum shorter than width of eye




pronotum with anterior margin moderately excavated



TROPICAL BED BUG
Cimex hemipterus
SO. U. S. & TROPICS


pronotum with anterior margin deeply excavated



BED BUG
Cimex lectularius
TEMPERATE AREAS


Fig. 139 KISSING BUGS: PICTORIAL KEY TO SOME COMMON SPECIES IN THE UNITED STATES

Harold George Scott and Margery R. Borom




STINGING CATERPILLARS:
PICTORIAL KEY TO SOME IMPORTANT UNITED STATES SPECIES
Harold George Scott & Chester J. Stojanovich

WITH DORSAL SADDLE



SADDLEBACK CATERPILLAR
Sibine stimulæ


BODY FUR-LIKE



PUSS CATERPILLAR
Megalopyge opercularis


WITHOUT DORSAL SADDLE

BODY NOT FUR-LIKE



7 PROLEGS


ABOUT 3/4-INCH LONG,
YELLOWISH GREEN




A SLUG CATERPILLAR
Euclea chloris

5 PROLEGS

ABOUT 2 INCHES LONG,
RED AND WHITE




IO MOTH CATERPILLAR
Automeris io



A FLANNEL MOTH CATERPILLAR
Ncraea cretata

MOTHS: KEY TO SOME SPECIES COMMONLY ASSOCIATED WITH STORED FOOD
Harold George Scott


- | | |
|---|--------------------------|
| 1. Caterpillars | 2 |
| Adult moths | 5 |
| 2. Pinkish larvae up to 3/5-inch long living in silken tubes and pro-
ducing matter webbing in the infested food (<i>Anagasta kuhniella</i>) | MEDITERRANEAN FLOUR MOTH |
| Whitish larvae with or without black or orange markings | 3 |
| 3. Black head and prothorax; orange markings at both ends of the body;
living in silken tubes (<i>Pyralis farinalis</i>) | MEAL MOTH |
| Without black head and prothorax | 4 |
| 4. White to greenish-white larvae producing matter webbing in the infested
food (<i>Plodia interpunctella</i>) | INDIAN MEAL MOTH |
| Whitish; not producing matted webbing; living inside kernels of grain
(<i>Sitotroga cerealella</i>) | ANGOUMOIS GRAIN MOTH |
| 5. Wings unicolorous to slightly spotted; long fringe at rear of wings
(<i>Sitotroga cerealella</i>) | ANGOUMOIS GRAIN MOTH |
| Wings heavily dark marked | 6 |
| 6. Distal half of front wings dark; basal half light (<i>Plodia inter-</i>
<i>punctella</i>) | INDIAN MEAL MOTH |
| Wings not so marked | 7 |
| 7. Basal and distal thirds of front wings dark; middle portion of front
wings light (<i>Pyralis farinalis</i>) | MEAL MOTH |
| Front wings pale gray with transverse wavy black markings
(<i>Anagasta kuhniella</i>) | MEDITERRANEAN FLOUR MOTH |



Angoumois Grain Moth


BLISTER BEETLES: KEY TO SOME COMMON UNITED STATES SPECIES

Harold George Scott and Chester J. Stojanovich



BEETLES: PICTORIAL KEY TO SOME SPECIES COMMONLY ASSOCIATED WITH STORED FOODS

Harry D. Pratt



STINGING HYMENOPTERA:
PICTORIAL KEY TO SOME COMMON UNITED STATES FAMILIES
Harold George Scott and Chester J. Stojanovich




DIAGRAM OF SOCIAL WASP





DIAGRAM OF FIRE ANT



HYMENOPTERA: KEY TO SOME COMMON SPECIES WHICH STING MAN
Harry D. Pratt and Chester J. Stojanovich

- | | |
|-------------------------------|-----|
| 1. With wings (Fig. 1 A)..... | 2 |
| Without wings (Fig. 1 B)..... | .32 |



Fig. 1 A




Fig. 1 B

- | | |
|--|-----|
| 2. First (and sometimes second) segment of the abdomen node-like, clearly separated above and below from rest of abdomen (Fig. 2 A). Nest in ground, wood, or buildings (Family Formicidae)..... | ANT |
|--|-----|

Abdomen with or without some constriction of first abdominal segments, but without true node formation of basal abdominal segments (Fig. 2 B)..... 3




Fig. 2 A




Fig. 2 B

- | | |
|---|---|
| 3. All hairs on body simple, unbranched; hind tarsus slender, first segment not broadened or thickened (Fig. 3 A). (Superfamilies Vespoidea and Sphecoidea). Wasps and Hornets..... | 4 |
|---|---|

At least some hairs on thorax branched or plumose; hind tarsus with first segment broadened and thickened, often densely hairy (Fig. 3 B). (Superfamily Apoidea). Bees..... 27




Fig. 3 A




Fig. 3 B

4. Pronotum extending entirely, or almost back, to the tegula (the scale covering base of fore-wing), its hind angles not lobed (Fig. 4 A). (Superfamily Vespoidea)..... 5

Pronotum shortened, more or less collar-like, not extending back to tegula, its hind angles often produced into lobes (Fig. 4 B). (Superfamily Sphecoidea)..... 22




Fig. 4 A




Fig. 4 B

5. Fore wing almost always folded when in repose; first discoidal cell very long, as a rule much longer than the submedian cell (Fig. 5 A). Both solitary and colonial species (Family Vespidae)..... 6

Fore wing very rarely folded; first discoidal cell shorter than submedian cell (Fig. 5 B). Solitary species..... 21




Fig. 5 A




Fig. 5 B

6. One spur at tip of middle tibia; claws bifid, split at tip (Fig. 6 A). (Subfamily Eumeninae)..... 18
Solitary Wasps.....

Two spurs at tip of middle tibia; claws tapering to point (Fig. 6 B)..... 7




Fig. 6 A




Fig. 6 B

7. Clypeus (upper lip) broadly truncate and more or less notched at apex (Fig. 7 A); hind wing without a lobe at anal angle (Fig. 7 B). (Subfamily Vespinae). Hornets, Yellow Jackets.....8
 Clypeus somewhat pointed at apex (Fig. 7 C); hind wing with a lobe at anal angle (Fig. 7 D). (Subfamily Polistinae). Paper Wasps.....15




Fig. 7 A




Fig. 7 C




Fig. 7 B




Fig. 7 D

8. Oculo-malar space long, more than half the length of next to last antennal segment; vertical carina on pronotum (Fig. 8 A).....9
 Oculo-malar space short, less than half the length of next to last antennal segment; no vertical carina on pronotum (Fig. 8 B).....11




Fig. 8 A




Fig. 8 B

9. Very large species, 20-30 mm. long, extensively reddish-brown; postocellar area of vertex at least as long as ocellar triangle in dorsal view (Fig. 9 A). Builds paper nest in homes or hollow trees. (*Vespa crabro germana*)..... GIANT HORNET

Smaller species, 8-20 mm. long; black species with white, ivory white, or yellowish markings; postocellar area of vertex not as long as ocellar triangle (Fig. 9 B)..... 10




Fig. 9 A





Fig. 9 B



10. Black and white species; first and second abdominal segments entirely black, second with very narrow pale markings at tip of first segments in some males (Fig. 10 A). Builds enclosed globular nests under eaves or in trees. (Vespa maculata).....BALD-FACED HORNET

Black and yellow species; yellowish posterior margins of first and second abdominal segments deeply notched (Fig. 10 B). Builds globular paper nests under eaves or in trees.....
(Vespa arenaria).....A YELLOW JACKET




Fig. 10 A




Fig. 10 B

11. Black and white species (Fig. 11 A). Builds paper nest in ground or on trees.....
(Vespa consobrina).....A HORNET

Black and yellowish species (Fig. 11 B). All build paper nests in ground.....12




Fig. 11 A




Fig. 11 B

12. Mesonotum with two, broad, longitudinal, curved yellowish stripes reaching almost from front to hind margins (Fig. 12 A). Eastern species (*Vespa squamosa*). California and Oregon species (*Vespa sulphurea*).....A YELLOW JACKET
 Mesonotum entirely black, or with two short yellowish stripes near scutellum (Fig. 12 B).....13




Fig. 12 A




Fig. 12 B

13. Yellowish postero-lateral margins of pronotum usually even, parallel-sided; clypeus with broad, dark, longitudinal stripe, often anchor-shaped (Fig. 13 A & B). Northern species.....
 (*Vespa vulgaris*).....A YELLOW JACKET

Yellowish postero-lateral margin of pronotum not parallel-sided; clypeus with short dark median stripe or one or more small dark spots (Fig. 13 C & D).....14




Fig. 13 A




Fig. 13 B




Fig. 13 C




Fig. 13 D

14. First antennal segment largely yellowish in front; eyes encircled by yellowish band on upper three-fourths (Fig. 14 A). Western species (Vespa pennsylvanica)..... A YELLOW JACKET
 First antennal segment largely or entirely blackish; eyes with a blackish area dorsally separating pale anterior and posterior orbital bands (Fig. 14 B). Eastern species (Vespa maculifrons).....
 A YELLOW JACKET




Fig. 14 A




Fig. 14 B

15. Body and all legs entirely or largely orange-colored (Fig. 15 A). Builds paper combs in walls of house or hollow trees. (Polistes rubiginosus)..... ORANGE PAPER WASP
 Body with some blackish markings; at least hind tarsi pale-colored (Fig. 15 B)..... 16




Fig. 15 A




Fig. 15 B

Abdomen yellowish, with blackish, yellowish, and reddish markings; mesonotum reddish; a yellowish band behind ocelli (Fig. 16 A). Builds single or double paper combs under eaves or in buildings. (*Polistes zebra*). ZEBRA PAPER WASP

Abdomen largely blackish, with one or more pale bands starting at posterior margin of first or second segment; mesonotum largely blackish; no yellowish band behind ocelli (Fig. 16 B) 17




Fig. 16 B

Fig. 16 A

17. Large species 20-25 mm. long, propodeum with coarse transverse striae (Fig. 17 A). Builds paper combs in bushes or trees. (*Polistes annularis*) LARGE PAPER WASP

Medium-sized species, 12-17 mm. long; propodeum with fine striae or essentially smooth (Fig. 17 B). Builds paper combs under eaves or in buildings. (*Polistes fuscatus pallipes*) DARK PAPER WASP




Fig. 17 A

Fig. 17 B

18. Slender species with extremely elongate first abdominal segment (Fig. 18 A). Builds small mud, potter nests provisioned with caterpillars. (Eumenes fraterna)..... POTTER WASP

Stocky species, with stout first abdominal segment (Fig. 18 B). Nest in holes in ground or wood, or old mud-dauber nests provisioned with caterpillars. (Odynerus species and Monobia species)..... SOLITARY WASPS




Fig. 18 A




Fig. 18 B

19. Mesopleuron divided by an oblique suture into upper and lower parts (Fig. 19 A). Usually nest in holes in ground provisioned with spiders or tarantulas (Family Psammocharidae)..... SPIDER AND TARANTULA WASPS

Mesopleuron not divided by such an oblique suture (Fig. 19 B)..... 20




Fig. 19 A



Fig. 19 B

20. Bases of middle and hind coxae not covered by plates (Fig. 20 A). Parasites of other wasps and bees nesting in ground..... VELVET ANTS
 Bases of middle, and sometimes hind, coxae covered by plates (Fig. 20 B)..... 21




Fig. 20 A




Fig. 20 B

21. Wing membrane beyond cells with wrinkles; inner margin of eye with a sinus; bases of middle and hind coxae covered by plates (Fig. 21 A & B). Male with three spines at tip of abdomen..... (Family Scoliidae)..... SCOLIID WASPS

Wing membrane beyond cells without wrinkles; inner margin of eye essentially straight; bases of middle coxae covered by plates (Fig. 21 C & D). Male with a single upturned spine at tip of abdomen. (Family Tiphidae)..... TIPHID WASPS




Fig. 21 B

Fig. 21 D




Fig. 21 A

Fig. 21 C

22. Very large species, 30 mm. long or more; first abdominal segment broad and sessile (Fig. 22 A)
Nest in holes in ground provisioned with cicadas. (*Sphecius speciosus*)..... CICADA KILLER

Smaller species, less than 25 mm. long; first abdominal segment longer and more slender (Fig.
22 B)..... 23




Fig. 22 A



Fig. 22 B

23. Eyes with deep sinus on inner side; one or two clearly defined submarginal cells; dark species
with whitish tarsus (Fig. 23 A). Builds organ-pipe mud nests. (*Trypoxyylon* species).....
PIPE ORGAN MUD-DAUBER

Eyes nearly straight on inner side; three well-defined submarginal cells; metallic blue, or
species with some pale markings on abdomen (Fig. 23 B & C)..... 24




Fig. 23 A




Fig. 23 B




Fig. 23 C

24. Petiole of abdomen two-segmented (Fig. 24 A). Nest in holes in ground. (Sphex species)
..... SOLITARY WASP

Petiole of abdomen one-segmented (Fig. 24 B)..... 25




Fig. 24 B

Fig. 24 A

25. Bright metallic-bluish species (Fig. 25 A). Builds mud nests provisioned with spiders.....
(Chalybion californicum)..... BLUE MUD-DAUBER

Darker species with yellowish or orange markings (Fig. 25 B)..... 26



Fig. 25 A

Fig. 25 B

26. Dark species with yellowish markings (Fig. 26 A). Builds mud nests provisioned with spiders.. .
(Sceliphron caementarium)..... COMMON MUD-DAUBER

Dark hairy species with orange markings (Fig. 26 B). Nest in holes in ground.
(Chlorion ichneumonea)..... ORANGE THREADED-WAISTED WASP



Fig. 26 A




Fig. 26 B

- Hind tibia without spurs (Fig. 27 A). Colony builds wax combs in bee hives, in houses, and in trees. (Apis mellifera)..... HONEY BEE
 Hind tibia with one or two spurs (Fig. 27 B)..... 28



Fig. 27 A




Fig. 27 B

28. Oculo-malar space longer than second segment of antenna; large hairy species with contrasting blackish and yellowish (sometimes reddish) pile (Fig. 28 A). Colony builds wax combs in nests in ground or logs, often in old mouse nests. (Family Bombidae; Bombus sp.).... BUMBLEBEES

Oculo-malar space short, eye reaching (or nearly reaching) base of mandible (Fig. 28 B).... 29




Fig. 28 A






Fig. 28 B

29. Very large species 15-25 mm. long with shiny bluish, nearly hairless upper abdomen; second submarginal cell strongly narrowed anteriorly (Fig. 29 A). Nest in holes bored in wood. (Xylocopa virginica) CARPENTER BEE
- Smaller species 2-14 mm. long, usually with some hairs on upper surface of abdomen, shiny greenish species; second submarginal cell not narrowed anteriorly (Fig. 29 B & C)..... 30



30. Fore-wing with two submarginal cells; abdomen of female with dense hairy patches on underside (Fig. 30 A). Builds nest out of leaves in tree holes (Megachile species) ... LEAFCUTTER BEES
- Fore-wing with three submarginal cells; abdomen without dense hairy patches on underside (Fig. 30 B & C)..... 31



- ... shiny greenish species (Fig. 31 A). Nest in ground. (Augochlora species).....
 METALLIC SOLITARY BEES
 ... duller species (Fig. 31 B). Nest in ground. (Halictus and Andrena species)....SOLITARY BEES



Fig. 31 A



Fig. 31 B

32. First (and sometimes second) segment of abdomen node-like (Fig. 32 A). Build colony nests in ground, under stones, in wood, or in buildings (Family Formicidae).....ANTS
 First and second segments of abdomen not node-like (Fig. 32 B)..... 33




Fig. 32 A



Fig. 32 B

33. Larger species 3-25 mm. long, usually with definite dark and reddish or orange-colored hairs (Fig. 33 A). Parasites of ground-nesting bees and wasps (Family Mutillidae). . . VELVET ANTS

Smaller species 1-2 mm. long, with few sparse hairs; body various shades yellowish to brownish (Fig. 33 B). Parasites of wood-boring beetles (Family Bethylidae, Scleroderma species... PARASITIC WASPS




Fig. 33 A





Fig. 33 B


ANTS: KEY TO SOME COMMON SPECIES
Harold George Scott

- | | |
|--|------------------------|
| 1. Pedicel ("waist") 1-segmented | 2 |
| Pedicel 2-segmented | 4 |
| 2. Petiole (scale on pedicel) poorly developed, hidden beneath abdomen
<i>(Tapinoma sessile)</i> | ODOROUS HOUSE ANT |
| Petiole well-developed, erect, not hidden beneath abdomen | 3 |
| 3. Tip of abdomen without circlet of hairs (<i>Iridomyrmex humilis</i>) | ARGENTINE ANT |
| Tip of abdomen with circlet of hairs (<i>Camponotus herculeanus pennsylvanicus</i>) | BLACK CARPENTER ANT |
| 4. Head and thorax with numerous spines (<i>Atta texana</i>) | TEXAS LEAF-CUTTING ANT |
| Head and thorax spineless or with 1 pair of spines on the posterior thorax | 5 |
| 5. Thorax and head covered with "fingerprints"; posterior thorax with
single pairs of spines (<i>Tetramorium caespitum</i>) | PAVEMENT ANT |
| Thorax and head without "fingerprints"; posterior thorax without spines | 6 |




- | | |
|---|-------------------|
| 6. Antennal club 2-segmented | 8 |
| Antennal club 3-segmented | 7 |
| 7. Shiny-black (<i>Monomorium minimum</i>) | LITTLE BLACK ANT |
| Yellowish-red (<i>Monomorium pharaonis</i>) | PHARAOH ANT |
| 8. House infesting ants (<i>Solenopsis molesta</i>) | THIEF ANT |
| Outdoor mound-building ants | 9 |
| 9. Mandibles strongly incurved (<i>Solenopsis geminata</i>) | TROPICAL FIRE ANT |
| Mandibles not strongly incurved | 10 |
| 10. Dorsal surface of head with large coarse, scattered punctures
<i>(Solenopsis saevissima var. richteri)</i> | IMPORTED FIRE ANT |
| Dorsal surface of head without punctures (<i>Solenopsis xyloni</i>) | SOUTHERN FIRE ANT |


DIPTERA: PICTORIAL KEY TO PRINCIPAL FAMILIES OF PUBLIC HEALTH IMPORTANCE
H. R. Dodge



DOMESTIC FLIES: PICTORIAL KEY TO COMMON SPECIES IN THE U.S.
H. R. Dodge




DOMESTIC FLIES: PICTORIAL KEY TO COMMON SPECIES
 Harold George Scott and Margery R. Borom



DOMESTIC FLIES: PICTORIAL KEY TO COMMON SPECIES IN SOUTHERN U.S.


H. R. Dodge



U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
Communicable Disease Center, Training Branch
Atlanta, Georgia – 1948, Revised Aug. 1953

V TO SOME COMMON SPECIES — J. M. Seago

FLY LARVAE: PICTORIAL



* ALL SPRAKLES REFERRED TO ARE LEFT SPRAKLES

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE, Communicable Disease Center, Training Branch, Atlanta, Georgia — 1952—Revised 1953

FLY LARVAE: KEY TO SOME SPECIES OF PUBLIC HEALTH IMPORTANCE
 Chester J. Stojanovich — Harry D. Pratt — Elwin E. Bennington

1. Larva with a definite, hard, sclerotized head capsule (Fig. 1 A)..... 2
- Larva without a definite, hard, sclerotized head capsule (Fig. 1 B)..... 3




Fig. 1 A




Fig. 1 B

2. Body flattened; large larvae 12-20 mm. long (Fig. 2 A)... (Hermetia illucens) SOLDIER FLY

Body cylindrical with spiracles opening in a tubular segment at posterior end of body, last segment modified into a sclerotized air tube (Fig. 2 B).....
 (Genus Psychoda & allies) FILTER FLIES




Fig. 2 A




Fig. 2 B

3. Body with spine-like dorsal and lateral processes on each segment; posterior spiracles on small elevations (Fig. 3 A)..... (Genus Fannia)... 4

Body smooth, or with short spines, but no long lateral processes (Fig. 3 B)..... 5




Fig. 3 A




Fig. 3 B

4. Processes branched or feathery (Fig. 4 A).....(Fannia scalaris) LATRINE FLY

Processes without branches, spiny (Fig. 4 B)..(Fannia canicularis) LESSER HOUSE FLY




Fig. 4 A




Fig. 4 B

5. Posterior spiracles on peg-like tubercles or cones; smaller larvae, usually 6-9 mm. long (Fig. 5 A)..... 6

Posterior spiracles not on peg-like tubercles; larger larvae, usually 9-18 mm. long (Fig. 5 B)..... 7




Fig. 5 A




Fig. 5 B

6. Posterior spiracles at ends of long tubercles (Fig. 6 A).....
.....(Genus Drosophila) VINEGAR FLIES

Posterior spiracles on short cones, last segment with short finger-like lateral process
(Fig. 6 B).....(Piophila casei) CHEESE SKIPPER




Fig. 6 A



Fig. 6 B

7. Posterior end of body extended to form a tail (Fig. 7 A).....
.....(Eristalis tenax) RAT-TAILED MAGGOT

Body swollen or tapered posteriorly, but never extended into a tail like process (Fig. 7 B).. 8




Fig. 7 A




Fig. 7 B

8. Peritreme present, with 3 distinct slits (Fig. 8 A)..... 9

Peritreme absent; or if present without 3 distinct slits (Fig. 8 B & C)..... 23




Fig. 8 A




Fig. 8 B




Fig. 8 C

9. Slits of posterior spiracles straight (Fig. 9 A)..... 10

Slits of posterior spiracles strongly sinuous (Fig. 9 B)..... 22




Fig. 9 A



Fig. 9 B

10. Dorsal and ventral arms of cephaloskeleton almost equal (Fig. 10 A); peritreme with two non-sclerotized areas away from the button (Fig. 10 B)..... (Genus Ophvra) DUMP FLY

Dorsal arm of cephaloskeleton longer than ventral arm (Fig. 10 C); peritreme complete or with one weakly sclerotized area (Fig. 10 D & E)..... 11




Fig. 10 A




Fig. 10 C




Fig. 10 B

Fig. 10 D




Fig. 10 E



11. Posterior spiracles with peritreme complete, sometimes weak in area of button (Fig. 11 A)..... 12

Posterior spiracles with peritreme incomplete, not enclosing a sometimes ill-defined button (Fig. 11 B)..... 16




Fig. 11 A




Fig. 11 B

12. Spiracular plate and button heavily sclerotized; accessory oral sclerite present (Fig. 12 A & B)..... 13

Spiracular plate and button not heavily sclerotized; accessory oral sclerite absent (Fig. 12 C & D)..... 14




Fig. 12 A




Fig. 12 B




Fig. 12 C





Fig. 12 D


13. Mandibular sclerite with tooth longer than greatest width of basal portion (Fig. 13 A).....
(Calliphora vicina) A BLUE BOTTLE FLY

Mandibular sclerite with tooth only as long as greatest width of basal portion (Fig. 13 B)..
(Cynomyopsis cadaverina) A BLUE BOTTLE FLY




14. Peritreme thick with rounded or sharp projections which extend inward toward spiracular slits (Fig. 14 A); cephaloskeleton as in figure 14 B.....
(Phaenicia caeruleiviridis) A GREEN BOTTLE FLY

Peritreme thin, usually with no projections or if present only slightly sclerotized (Fig. 14 C).....15



15. At least one of the prothoracic spiracles with 8 or more openings (Fig. 15 A); peritreme and cephaloskeleton as in figures 15 B & C..
(Phaenicia sericata) A GREEN BOTTLE FLY

At least one of the prothoracic spiracles with 6 or less openings (Fig. 15 D); peritreme and cephaloskeleton as in figures 15 E & F.....
(Syn. *P. pallescens*).....
(Phaenicia cuprina) A BRONZE BOTTLE FLY



16. Spiracular slits not pointing toward opening in peritreme (Fig. 16 A).....17
 Spiracular slits pointing toward opening in peritreme (Fig. 16 B).....18




Fig. 16 A




Fig. 16 B

17. Very large size, about 20 mm. long; mandibular sclerite as in figure 17 A.....
(Sarcophaga clitellivora or S. bullata) A FLESH FLY
 Smaller size, about 10 mm. long; mandibular sclerite as in figure 17 B.....
(Sarcophaga haemorrhoidalis) A FLESH FLY



Fig. 17 A



Fig. 17 B

18. At least one of the prothoracic spiracles with 9 or less openings (Fig. 18 A).....19
 At least one of the prothoracic spiracles with 10 or more openings (Fig. 18 B).....20



Fig. 18 A



Fig. 18 B

19. Mandibular sclerite with tooth longer than width of basal portion (Fig. 19 A).....
(Wohlfahrtia opaca) A FLESH FLY
 Mandibular sclerite with tooth only as long as greatest width of basal portion (Fig. 19 B)..
(Wohlfahrtia vigil) A FLESH FLY




Fig. 19 A




Fig. 19 B

20. Button is distinct or absent; walls of slits with lateral swellings (Fig. 20 A)..... 21
 Button present, walls of slits without lateral swellings (Fig. 20 B).....
 Phormia regina BLACK BLOW FLY



Fig. 20 A




Fig. 20 B

21. Tracheal trunks pigmented (Fig. 21 A).....
 (Cochliomyia hominivorax) PRIMARY SCREW-WORM
 Tracheal trunks not pigmented (Fig. 21 B).....
 (Cochliomyia macellaria) SECONDARY SCREW-WORM




Fig. 21 A




Fig. 21 B

22. Peritreme thick (Fig. 22 A)..... (Musca domestica) HOUSE FLY
 Peritreme thin (Fig. 22 B)..... (Haematobia irritans) HORN FLY




Fig. 22 A




Fig. 22 B

23. Small or slender, round larvae, usually less than 13 mm. long, tapering anteriorly (Fig. 23 A)..... 24
 Large, robust larvae, over 15 mm long, with very stout spines (Fig. 23 B)..... 26



Fig. 23 A




Fig. 23 B

24. Button centrally located (Fig. 24 A)..... (Stomoxys calcitrans) STABLE FLY
 Button not centrally located (Fig. 24 B)..... 25




Fig. 24 A




Fig. 24 B

25. Slits of posterior spiracles strongly sinuous (Fig. 25 A).... (Musca autumnalis) FACE FLY
 Slits of posterior spiracles not strongly sinuous (Fig. 25 B)..... (Genus Mucina) FALSE STABLE FLY



Fig. 25 A




Fig. 25 B

26. Posterior spiracles with 3 distinct slits (Fig. 26 A)..... 27
 Posterior spiracles without 3 distinct slits (Fig. 26 B)..... 28




Fig. 26 A




Fig. 26 B

27. Spiracular slits straight and sunken in deep cavity (Fig. 27 A); body shape as in figure 27 B. (Genus Dermatobia) HUMAN BOT FLY

Spiracular slits curved and at most in shallow cavity (Fig. 27 C); body shape as in figure 27 D. (Genus Gasterophilus) HORSE BOT FLY




Fig. 27 A




Fig. 27 C




Fig. 27 B




Fig. 27 D

28. Each spiracle divided into several plates (Fig. 28 A).....
..... (Genus Cuterebra) RABBIT AND RODENT BOT FLY

Each spiracle not divided into several plates (Fig. 28 B)..... 29




Fig. 28 A




Fig. 28 B

29. Button centrally located (Fig. 29 A)..... (Oestrus ovis) SHEEP BOT FLY

Button not centrally located (Fig. 29 B)..... 30




Fig. 29 A




Fig. 29 B

30. Opening toward button narrow (Fig. 30 A).... (Hypoderma bovis) NORTHERN CATTLE GRUB

Opening toward button wide (Fig. 30 B)..... (Hypoderma lineatum) CATTLE GRUB



Fig. 30 A





Fig. 30 B


MOSQUITOES: CHARACTERISTICS OF ANOPHELINES AND CULICINES

Kent S. Littig and Chester J. Stojanovich


ANOPHELES




AEDES




CULEX




AEDES AEGYPTI
Chester J. Stojanovich



ANOPHELES QUADRIMACULATUS
Harry D. Pratt




CULEX QUINQUEFASCIATUS
Harry D. Pratt



MOSQUITOES: PICTORIAL KEY TO U.S. GENERA OF LARVAE


Harry D. Pratt



MOSQUITOES: PICTORIAL KEY TO SOME COMMON LARVAE OF THE UNITED STATES

139•


Chester J. Stojanovich and Harry D. Pratt



MOSQUITOES: PICTORIAL KEY TO SOME COMMON LARVAE OF WESTERN UNITED STATES


Harry D. Pratt

• 140




U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE, PUBLIC HEALTH SERVICE, Communicable Disease Center, Training Branch, Atlanta, Georgia - 1960


MOSQUITOES: PICTORIAL KEY TO SOME LARVAE
COMMONLY FOUND IN ARTIFICIAL CONTAINERS
Harry D. Pratt and Chester J. Stojanovich



MOSQUITOES: PICTORIAL KEY TO ANOPHELINE LARVAE OF THE UNITED STATES
Stanley B. Freeborn and Eugene J. Gerberg




MOSQUITOES: PICTORIAL KEY TO SOME LARVAE OF FLORIDA
COMMONLY FOUND IN CONTAINERS
Chester J. Stojanovich and Harry D. Pratt



PART II


air tube with one hair or tuft on each side



air tube with several hairs or tufts on each side




comb scales with a strong median spine,
thorax with prominent lateral spines



Aedes aegypti

comb scales with a fringe, thorax
with less prominent lateral spines




Aedes triseriatus


air tube at least 6 times as long as wide



air tube 3-5 times as long as wide




comb scales in a single row
air tube with a basal tuft




Culiseta melanura


comb scales in a patch,
air tube without a basal tuft




antenna with tuft
inserted at middle
air tube with 3 pairs
of long single hairs



antenna with tuft
inserted beyond middle
air tube with 4 pairs
of multiple tufts




thorax with spicules,
lateral hair of saddle single



Culex nigripalpus

thorax without spicules,
lateral hair of saddle double

thorax without spicules,
lateral hair of saddle double




Culex salinarius

Culex restuans


Culex quinquefasciatus

MOSQUITOES: PICTORIAL KEY TO SOME COMMON LARVAE OF PUERTO RICO
FOUND IN CONTAINERS


Harry D. Pratt and Chester J. Stojanovich




MOSQUITO DIAGRAM - ADULT FEMALE AEDES
Chester J. Stojanovich and Harold George Scott



MOSQUITO DIAGRAM – LATERAL ASPECT OF MOSQUITO THORAX
Chester J. Stojanovich



MOSQUITOES: PICTORIAL KEY TO UNITED STATES GENERA OF ADULTS (FEMALE)
Harry D. Pratt and Chester J. Stojanovich



proboscis stout on basal half, outer half tapered and strongly turned downward



abdominal scales dark dorsally and pale ventrally; postnotum with setae



wing with second marginal cell less than half as long as its petiole



abdomen blunt



base of subcosta with row of bristles on under side




base of subcosta without row of bristles on under side



dorsal segments of abdomen with pale scales apically, or if absent, hind tibia with long, erect scales




dorsal segments of abdomen with pale scales basally, hind tibia without erect scales




most wing scales at base of wing narrow, dark



antenna not longer than proboscis, first flagellar segment about as long as following segments



antenna much longer than proboscis, first flagellar segment as long as next two segments



mesonotum with fine longitudinal lines of white scales




mesonotum without lines of white scales




MOSQUITOES: PICTORIAL KEY TO SOME COMMON ADULTS (FEMALE) OF THE UNITED STATES

Harry D. Pratt and Chester J. Stojanovich




MOSQUITOES: PICTORIAL KEY TO SOME ADULTS (FEMALE) IN EASTERN UNITED STATES

Harry D. Pratt and Chester J. Stojanovich




MOSQUITOES: PICTORIAL KEY TO SOME COMMON ADULTS (FEMALE) OF WESTERN UNITED STATES

Harry D. Pratt and Chester J. Stojanovich




MOSQUITOES: PICTORIAL KEY TO SOME ADULTS
COMMONLY ASSOCIATED WITH AEDES AEGYPTI
Harry D. Pratt and Chester J. Stojanovich




MOSQUITOES: PICTORIAL KEY TO ADULT FEMALE ANOPHELINES OF UNITED STATES

Richard H. Daggy




MOSQUITOES: PICTORIAL KEY TO UNITED STATES GENERA
BASED ON MALE GENITALIA
PART I
Chester J. Stojanovich

basistyle about equal in length to dististyle
and with 1-2 stout setae near base





Anopheles

basistyle usually much shorter than dististyle,
without 1-2 stout setae near base




claspette present 'absent only in *Aedes aegypti* as shown below'




Aedes aegypti

claspette absent




SEE PART II SECTION II

distal lobe of basistyle with leaf-like scales



Haemagogus


distal lobe when present without leaf-like scales




SEE PART II SECTION I

PART II**PART II SECTION I**

dististyle not slender but variously shaped as shown below





dististyle slender 'exceptions being
Aedes cinereus with dististyle furcate at base and
Aedes vexans with claw of dististyle not inserted at tip

***Psorophora******Aedes*****PART II SECTION II**



tenth sternite crowned with teeth or tuft of spines

tenth sternite simple or with few teeth

**SEE PART III**


distal lobe of basistyle with leaf-like scales or rods
 lobe of ninth tergite short

distal lobe of basistyle without leaf-like scales or rods
 lobe of ninth tergite half as long as basistyle




PART III

dististyle greatly modified at apex




Wyeomyia

dististyle not greatly modified at apex




claw of dististyle comb-like




Orthopodomyia

claw of dististyle not comb-like




SEE PART IV


PART IV




MOSQUITOES: PICTORIAL KEY TO MOST ADULTS (FEMALE) OF NEW MEXICO
PART I
Chester J. Stojanovich




PART II




MOSQUITOES: PICTORIAL KEY TO SOME COMMON ADULTS (FEMALE) OF IDAHO
Chester J. Stojanovich




MOSQUITOES: PICTORIAL KEY TO ALL ADULTS (FEMALE) OF TEXAS
PART I
Chester J. Stojanovich




PART II




PART III




PART IV




MOSQUITOES: PICTORIAL KEY TO SOME COMMON ADULTS (FEMALE) OF UTAH
PART I
 Chester J. Stojanovich




PART II



CAT FLEA - CTENOCEPHALIDES FELIS
adult female




FLEA DIAGRAM — WITH STRUCTURES LABELED
Harry D. Pratt



FLEAS: PICTORIAL KEY TO SOME COMMON SPECIES IN THE UNITED STATES

Harry D. Pratt



FLEAS: PICTORIAL KEY TO SPECIES FOUND ON DOMESTIC RATS IN SOUTHERN UNITED STATES

Slightly revised April 1947
By Roy F Fritz and Harry D Pratt
Rev March, 1954

Genal and pronotal
combs present




Genal, pronotal and abdominal combs present

No combs



Pronotal comb present



FLEAS: ILLUSTRATED KEY TO SPECIES FOUND DURING PLAGUE INVESTIGATIONS
Harry D. Pratt and Chester J. Stojanovich

1. Pronotal and genal combs absent (Fig. 1 A).....2

Pronotal combs present; genal comb present or absent (Fig. 1 B & C)...5




Fig. 1 A

pronotal comb

Fig. 1 B

genal comb

Fig. 1 C

2. Front margin of head with two angles; three thoracic tergites together shorter than the first abdominal tergite (Fig. 2 A). (Echidnophaga gallinacea).....STICKTIGHT FLEA

Front margin of head rounded; three thoracic tergites together longer than the first abdominal tergite (Fig. 2 B).....3




Fig. 2 A

Fig. 2 B

3. Ocular bristle in front of eye; mesopleuron divided by internal sclerotization; female with spermatheca partially pigmented (Fig. 3 A & B)...
.....(Genus Xenopsylla).....4

Ocular bristle beneath eye; mesopleuron without internal sclerotization; female with spermatheca entirely without pigment (Fig. 3 C & D).....
(Pulex irritans).....HUMAN FLEA





Fig. 3 A

Fig. 3 B


Fig. 3 C

Fig. 3 D

4. Genus Xenopsylla



Xenopsylla cheopis, male terminal segments.



Xenopsylla cheopis, female terminal segments

X. vexabilis hawaiiensis

5. Genal comb absent (Fig. 5 A).....6

Genal comb present (Fig. 5 B).....8




Fig. 5 A




Fig. 5 B

genal comb

6. Pronotal comb with about 12 teeth on each side (Fig. 6 A). India.....
.....*Stivalius ahalaee*

Pronotal comb with 5 to 10 teeth on each side (Fig. 6 B).....7




Fig. 6 A




Fig. 6 B

7. Labial palpus long, extending beyond trochanter of first leg (Fig. 7 A).
Diamanus montanus.....ROCK SQUIRREL FLEA

Labial palpus short, not extending to tip of coxa of first leg (Fig. 7 B).
Nosopsyllus fasciatus.....NORTHERN RAT FLEA




Fig. 7 A

trochanter-----

labial palpus-----




Fig. 7 B

labial palpus-----

trochanter-----

3. Genal comb with two teeth (Fig. 8 A).....(Genus Neopsylla)
Neopsylla setosa important in U. S. S. R., Mongolia and Manchuria.

Genal comb with three teeth (Fig. 8 B).....(Genus Ctenophthalmus)
Ctenophthalmus breviusculus and pollex potential vectors in U. S. S. R.

Genal comb with four teeth (Fig. 8 C).....(Genus Leptopsylla)
Leptopsylla segnis is cosmopolitan.




Fig. 8 A




Fig. 8 B




Fig. 8 C

Genal comb with more than five teeth.....(Genus Ctenocephalides). 9

9. Head strongly rounded anteriorly; first spine of genal comb about half as long as second; hind tibia with the spiniform setae A and B (Fig. 9 A & B).
Ctenocephalides canis.....DOG FLEA

Head not strongly convex anteriorly; first spine of genal comb almost as long as second spine; hind tibia with spiniform seta B, spiniform seta A replaced by a minute seta which may be absent in some specimens (Fig. 9 C & D). Ctenocephalides felis.....CAT FLEA




Fig. 9 A




Fig. 9 C




Fig. 9 B





Fig. 9 D

SNAKES: PICTORIAL KEY TO VENOMOUS SPECIES IN UNITED STATES

PART I



Chester J. Stojanovich and Margaret A. Parsons



PART II



head with large scales medially

head with small scales medially







upper preocular usually touching postnasal

upper preocular and postnasal separated


*Sistrurus catenatus*
MASSASAUGA RATTLESNAKE*S. c. catenatus*
Great Lakes & Central U.S.*S. c. edwardsii*
Arizona, Colorado, New Mexico, Texas*S. c. tergeminus*
Colorado, Kansas, Nebraska, New Mexico, Oklahoma, Texas*Sistrurus miliaris*
PIGMY RATTLESNAKE*S. m. miliaris*
Southeastern*S. m. barbouri*
Southeastern*S. m. streckeri*
Southeastern

supraocular scale modified into a hornlike ridge

supraocular scale not modified into a hornlike ridge


*Crotalus cerastes*
SIDEWINDER RATTLESNAKE*C. c. cerastes*
Arizona, California, Nevada, Utah*C. c. cercobombus*
Arizona*C. c. laterorepens*
Arizona, California*Crotalus willardi*
RIDGE-NOSED RATTLESNAKE*C. w. silus*
New Mexico*C. w. willardi*
Arizonadorsal blotches on body divided into 2 parallel rows
first supralabial scale broadly attached to postnasal scale*Crotalus pricei*

TWINSPOTTED RATTLESNAKE Arizona


dorsal blotches on body not divided into 2 parallel rows
first supralabial scale not broadly attached to postnasal scale, sometimes completely separated

SEE PART III


PART III



PART IV




**BIRDS: PICTORIAL KEY TO SOME COMMON PEST SPECIES
OF PUBLIC HEALTH IMPORTANCE**




**U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE, Communicable Disease Center, Training Branch, Atlanta, Georgia - 1964**

PIGEON, *COLUMBA LIVIA* — EXTERNAL MORPHOLOGY
Harold George Scott and Walter S. Dougherty




RODENTS: PICTORIAL KEY TO SOME COMMON UNITED STATES GENERA

Harold George Scott and Margery R. Borom




DOMESTIC RODENT FIELD IDENTIFICATION
Robert Z. Brown

ROOF RAT *Rattus rattus*




NORWAY RAT *Rattus norvegicus*



SCALE IN INCHES

A horizontal scale bar marked at 0, 1, 2, and 3 inches.

DOMESTIC RODENTS AND COCKROACHES: PICTORIAL KEY TO DROPPINGS
Harold George Scott and Margery R. Borom




*All characteristics for average, dry, adult droppings. Study groups, not individual droppings.


PRAIRIE DOGS: PICTORIAL KEY TO COMMON NORTH AMERICAN SPECIES
(*Cynomys*)

Harry Weinburgh and Margery R. Borom

Tail white-tipped, long, more than 1/3 total length (72-115 mm.)



Tail white-tipped, short, less than 1/3 total length (40-58 mm.)



Black on tail covering most of distal half
MEXICO ONLY

Black on tail confined to distal third
TEXAS TO SASKATCHEWAN



MEXICAN PRAIRIE DOG
C. mexicanus



BLACK-TAILED PRAIRIE DOG
C. ludovicianus



Terminal half tail white without dark center




GUNNISON PRAIRIE DOG
C. gunnisoni

Terminal half tail with dark center (gray)



Summer color reddish (cinnamon or clay color mixed with buff); darker on rump
CENTRAL VALLEYS OF UTAH




UTAH PRAIRIE DOG
C. parvidens

Summer color grayish (pinkish buff mixed with black); dark patch on cheek and above eye
WYOMING, COLORADO, AND EASTERN UTAH




WHITE-TAILED PRAIRIE DOG
C. leucurus

RABBITS AND HARES: PICTORIAL KEY TO COMMON UNITED STATES SPECIES
 Harold George Scott and Margery R. Borom



BATS: PICTORIAL KEY TO UNITED STATES GENERA
Harold George Scott and Chester J. Stajanovich



U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE, Communicable Disease Center, Training Branch, Atlanta, Georgia - 1960 - Revised 1962

SELECTED REFERENCES

GENERAL

Mallis A. 1964. Handbook of pest control. MacNair Dorland Co., New York, 1148 pp.

Metcalf, C.L., Flint, W.P., and Metcalf, R.L. 1962. Destructive and useful insects. McGraw-Hill Book Co., Inc., New York, xii + 1087 pp.

CRUSTACEA

Barnes, R.D. 1963. Invertebrate zoology. W.B. Saunders Co., Philadelphia, pp. 380-474.

Pennak, R.W. 1953. Fresh water invertebrate of the United States. Ronald Press, New York, pp. 321-469.

CENTIPEDES

Cloudsley-Thomson, J. 1958. Spiders, scorpions, centipedes and mites. Permagon Press, New York, xii + 228 pp.

Minton, S. 1959. Centipedes. Pest Control, 27(3):29.

MILLIPEDES

Chamberlain, R., and Hoffman, R. 1958. A check list of the millipedes of North America. U.S. Natl. Mus. Bull. 212, 236 pp.

Halstead, B., and Ryckman, R. 1949. Injurious effects from contacts with millipedes. Med. Arts. Sci., 3:16-18.

SPIDERS

Emerton, J.H. 1961. The common spiders of the United States. Dover Publications, Inc., New York City, xx + 227 pp.

Kaston, B.J., and Kaston, E. 1953. How to know the spiders. Wm. C. Brown Co., Dubuque, vi + 22 pp.

SCORPIONS

Ewing, H.E. 1928. The scorpions of the western part of the United States with notes on those occurring in Northern Mexico. Proc. U.S. Natl. Mus., 73:1-24.

Stahnke, H.L. 1956. Scorpions. Poisonous Animals Research Lab., Arizona State University, Tempe, 40 pp.

TICKS

Cooley, R.A., and Kohls, G.M. 1944. The Argasidae of North America, Central America, and Cuba. Amer. Midland Nat., Mono. #1, 152 pp.

Gregson, J.D. 1956. The Ixodoidea of Canada, Canada Dept. Agr., Sci. Serv., Publ. 930, 92 pp.

MITES

Baker, E.W., Evans, T.M., Gould, D.J., Hull, W.B., and Keegan, H.L. 1956. A manual of parasitic mites of medical or economic importance. Natl. Pest Control Assoc. Tech. Publ., New York, 170 pp.

Hughes, A.M. 1961. The mites of stored food. H.M. Stationery Off., London, England, vi + 287 pp.

SILVERFISH

Slabaugh, R.E. 1940. A new thysanuran, and a key to the domestic species of Lepismatidae (Thysanura) found in the U.S. Ent. News, 51:95-98.

Womersley, H. 1939. Primitive insects of South Australia. Handbook Flora Fauna So. Aust., Adelaide (Government of So. Aust.), 322 pp.

COLLEMBOLA

Scott, H. G. 1966. Insect Pests - Part I: Springtails. Mod. Maintenance Management, 18(9):19-21.

Scott, H.G. 1962. Collembola infesting man. Ann. Ent. Soc. America, 55(4): 428-430.

COCKROACHES

Roth, L.M., and Willis, E.R. 1957. The medical and veterinary importance of cockroaches. Smithsonian Misc. Coll., 134(10): V + 147 pp.

Willis, E.R., Riser, G.R., and Roth, L.M. 1958. Observations on reproduction and development in cockroaches. Ann. Ent. Soc. Amer., 51(1):53-69.

TERMITES

Snyder, T.C. 1954. Order Isoptera: the termites of the United States and Canada. Natl. Pest Control Assoc., Tech. Bull., 64 pp.

Scott, H.G. 1961. Keys to common North American domestic termites. Pest Control, 29(9):46.

EARWIGS

Hebard, M. 1934. The Dermaptera and Orthoptera of Illinois. Illinois Natl. Hist. Surv. Bull., 29(3):125-279.

Townes, H.K. 1945. A list of the generic and subgeneric names of Dermaptera with their genotypes. Ann. Ent. Soc. America, 38:343-356.

PSOCIDS

Broadhead, E. 1950. A revision of the genus Liposcelis . . . Trans. Roy. Ent. Soc. London, 101:335-338.

LICE (*Anoplura*)

Buxton, P.A. 1946. The louse. Williams & Wilkins Co., Baltimore, viii + 164 pp., 47 figs.

Pratt, H.D., and Littig, K.S. 1961. Lice of public health importance and their control. U.S. Govt. Print. Office, Washington, 16 pp.

LICE (*Mallophaga*)

Emerson, K.C. 1962. A tentative list of Mallophaga for (1) North American Birds (II) North American Mammals. Dugway Proving Grounds, Dugway, Utah, 217 + 20 pp.

LICE (Mallophaga) (continued)

Hopkins, G.H.E., and Clay, T. 1952. A check list of the genera and species of Mallophaga. British Mus. (Natl. Hist.), iii + 362 pp.

BUGS

Smith, C.N. 1958. Control of bed bugs. Pest Control, 26(11):9-12.

Usinger, R.L. 1944. The Triatominae of North and Central America and the West Indies and their public health significance. U.S. Public Health Service Bull. 288, iv + 83 pp.

LEPIDOPTERA

Allard, H.F., and Allard, H.A. 1959. Venomous moths and butterflies. J. Washington Acad. Sci., 48:18-21.

Goldman, L., Sawyer, F., Levine, A., Goldman, J., Goldman, S., and Spinanger, J. 1960. Investigative studies of skin irritations from caterpillars. J. Invest. Dermat., 34:67-79.

BEETLES

Cotton, R.T. 1956. Pests of stored grain and grain products. Burgess Publ. Co., Minneapolis, iii + 306 pp.

Pratt, H.D., and Scott, H.G. 1962. A key to some beetles commonly found in stored foods. Proc. Ent. Soc. Washington, 64(1):43-50.

HYMENOPTERA

Creighton, W.S. 1950. The ants of North America. Mus. Comp. Zool. Bull., 104:1-585.

Muesebeck, C.F.W., Krombein, K.V., Townes, H.K. 1951 and 1958. Hymenoptera of America North of Mexico + First Supplement. U.S. Dept. Agr. Monograph 2:1420 pp.

FLIES

- Stone, A., Sabrosky, C. W., Wirth, W. W., Foote, R. H., and Coulson, J. R. 1965. A catalog of the diptera of America north of Mexico. Agri. Handbook No. 276, U. S. Govt. Print. Off., Washington, D. C.
- Hall, D. 1948. The blowflies of North America. Thomas Say Foundation. Columbus, 477 pp.

MOSQUITOES

- Carpenter, S.J., and La Casse, W.J. 1955. Mosquitoes of North America. Univ. California Press, Berkeley, vi + 360 pp.

- Stojanovich, C.J. 1960-61. Illustrated key to common mosquitoes of (I) South-eastern United States, (II) Northeastern North America. Publ. by Author, P.O. Box 727, Emory University Branch, Atlanta, Georgia, 36 + 49 pp.

FLEAS

- Holland, G.P. 1949. The Siphonaptera of Canada. Canada Dept. Agr., Publ. 817, Tech. Bull. 70, 306 pp.

- Stark, H.E. 1958. The Siphonaptera of Utah. U.S. Dept. Health, Education, and Welfare, PHS, Atlanta, Georgia, xiii + 239 pp.

SNAKES

- Ditmars, R.L. 1939. A field book of North American snakes. Doubleday and Co., Inc., New York, xii + 305 pp. + 49 plates.

- Schmidt, K.P., and Davis, D.D. 1941. Field book of snakes of the United States and Canada. G.P. Putnam's Sons, New York, xiii + 365 pp.

BIRDS

- Scott, H.G. 1961. Pigeons: Public health importance and control. Pest Control, 29(9):9-20 + 60-61.

- Scott, H.G. 1964. Pigeon-borne disease control through sanitation and pigeon stoppage. Pest Control, 32(9):14-19 + 38-44.

RODENTS, RABBITS, AND HARES

Bjornson, B.F., and Wright, Charles V. 1960. Control of domestic rats and mice. U.S. Govt. printing Office, Washington,D.C., PHS Publ. 563, 25 pp.

Hall, E.R., and Kelson, K.R. 1959. The mammals of North America. The Ronald Press Co., New York, Vol. I, xxx + 625 pp., Vol. II, viii + 1162 pp.

BATS

Allen, G.M. 1939. Bats. Dover Publ. Inc., New York, x + 368 pp.

Scott, H.G. 1961. Bats: public health importance, identification and control. Pest Control, 29:23-26.